

## 1

## Clinical Pharmacology

*Leigh A. Lamont and Kurt A. Grimm*

### Introduction

Pharmacology is the study of drugs and their interactions with organisms (Page & Maddison 2002). Pharmacology incorporates aspects of statistics, biochemistry, biology, pathology, and medicine. Failure to interpret the description of drugs' pharmacological properties in the context of the clinical picture (i.e., clinical pharmacology) can result in unintended outcomes.

The pharmacological data available for most drugs are mean values derived from a relatively small number of individuals (usually healthy individuals). While this approach provides a starting point for clinical use of drugs, individual responses can vary greatly due to disease states, body condition, environment, genetics, coadministered drugs, and many other factors. When the toxic dose is close to the therapeutic dose (as is often the case with drugs used for immobilization and anesthesia), careful titration of dose and patient monitoring are required. However, the nature of working with wildlife and captive nondomestic species often precludes baseline health assessment, individualization of dosing, and intensive patient monitoring. This is one factor associated with increased risk of adverse outcomes when capturing or anesthetizing nondomestic species. It should also be appreciated that advances in drug safety will likely result in only limited improvement of the safety of anesthesia and immobilization. Management of other risk factors through airway management, reduction of stress, and improvements in supportive care will also be beneficial.

### Pharmacokinetics

Pharmacokinetics can be generally defined as what an organism does to a drug. Absorption, distribution, biotransformation, and elimination are processes which

determine the concentration of drug at the site of action (i.e., biophase). Pharmacokinetic parameters are estimates of these processes in the group of animals studied. These estimates can be used to predict or understand the way a drug interacts with an organism. It is important to understand that pharmacokinetic parameters can vary between individual animals and can be influenced by many different drug- and organism-related factors. Additionally, pharmacokinetic parameters are derived using mathematical models selected by the investigator. There is usually no correlation between model components and anatomical structures.

### Pharmacodynamics

Pharmacodynamics can be generally defined as what a drug does to an organism. Pharmacodynamics includes intended drug effects as well as adverse drug actions. Drugs such as opioids, alpha-2 adrenergic agonists, and antimuscarinics act by binding to relatively well-characterized receptor complexes located on cellular membranes and produce well-characterized effects. Nonsteroidal anti-inflammatory drugs (NSAIDs) inhibit prostaglandin production by binding to cyclooxygenase enzyme isoforms. Relating plasma drug concentrations to observed NSAID actions can be complex in comparison to other drugs (e.g., opioids) due to the different nature of the action. Preexisting prostaglandins, as well as the slower process of inhibiting an enzyme system, confound the relationship between drug concentration and effect. The molecular and neurophysiologic actions of inhalant anesthetics, in contrast, have not been completely characterized even though the drugs have been used clinically for many years and their pharmacodynamic effects have been well described (Steffey et al. 2015).

Pharmacodynamic effects are predictable for most clinically used drugs. However, individual animal responses can vary considerably. Additionally, the nature of capture of free-ranging and captive wildlife often makes accurate dosing and drug delivery difficult or impossible. Therefore, close monitoring of patient response and preparation for supportive care are paramount to safe immobilization and anesthesia.

Inhalant Anesthetics

Inhalant anesthetics are commonly used in companion animal veterinary practice. Their use under field conditions is limited due to the requirement for specialized delivery devices and a supply of delivery gas (e.g., oxygen). However, inhalant anesthetics are used commonly in controlled settings such as zoological parks and research laboratories because of the ease of titration of anesthetic depth and rapidity of recovery. Inhalant anesthetics should be delivered by a well-maintained anesthetic machine and properly trained individuals. While inhalant anesthetics are relatively safe, their low therapeutic index mandates frequent and careful monitoring of anesthetic depth.

Physics of Gases and Vapors

An understanding of the processes which influence the uptake and delivery of inhalant anesthetics allows the anesthesiologist to predict and respond to individual circumstances.

Brief Review of Molecular Theory

Molecules in a liquid state have more vibrational energy than when in a solid state, and each molecule can move through the liquid. If heat is added to a liquid, each molecule gains more kinetic energy and eventually some overcome the forces exerted by their neighbors and escape into the space above the liquid. This state is that of a gas or a vapor. A gas is a phase of matter that expands indefinitely to fill a containment vessel. A vapor is the gaseous state of a material below its boiling point.

A vapor is in equilibrium with the liquid beneath it. Because both gaseous and liquid molecules have kinetic energy, they are in constant motion. The molecules in the vapor phase are striking the liquid–gas interface and returning to the liquid, while liquid molecules are leaving the interface to become vapor. The relationship between these two phases depends mainly on the physicochemical properties of the molecules and the temperature of the system.

Vapor Pressure

Molecules in a gaseous state possess kinetic energy and collide with the walls of the containment vessel. These collisions produce a force on the walls. This force is spread over a surface area and therefore is a pressure (pressure = force/area). This pressure is called the vapor pressure. Since kinetic energy increases directly with temperature, vapor pressure must always be given with reference to the temperature it was measured at (e.g., vapor pressure of water is 47 mmHg at 37°C).

When many gases are present in a mixture, such as with atmospheric air or during delivery of inhalant anesthetics, each gas has a vapor pressure that is independent of the other gases (Dalton’s law of partial pressures). It is convention to refer to vapor pressure as partial pressure under these conditions. Partial pressure of an anesthetic agent is analogous to the concept of “free drug” and is important for determining the effect of the anesthetic (e.g., the level of central nervous system (CNS) depression correlates directly with the partial pressure of isoflurane within the brain) (see Table 1.1) (Steffey et al. 2015).

Vapor Concentration

Vapor (i.e., partial) pressure is important for the observed pharmacological effect of inhalant anesthetics. However, almost all anesthesiologists refer to the amount of anesthetic delivered in units of volumes % (said as volumes-percent), or just percent, which is a concentration. The fundamental difference between anesthetic partial pressure and anesthetic concentration is that partial pressure relates to the absolute number of molecules and their kinetic energy whereas concentration refers to the number of molecules of anesthetic relative to the total number of molecules present.

Table 1.1 Anesthetic agent vapor pressures at 20 and 24°C.

| Anesthetic Agent | Vapor Pressure @ 20°C in mmHg | Vapor Pressure @ 24°C in mmHg |
|------------------|-------------------------------|-------------------------------|
| Methoxyflurane   | 23                            | 28                            |
| Sevoflurane      | 160                           | 183                           |
| Enflurane        | 172                           | 207                           |
| Isoflurane       | 240                           | 286                           |
| Halothane        | 243                           | 288                           |
| Nitrous oxide    | Gas                           | Gas                           |

Source: Adapted from Steffey EP, Mama KM, Brosnan RJ. 2015. Inhalation anesthetics. In: *Veterinary Anesthesia and Analgesia: The Fifth Edition of Lumb and Jones* (KA Grimm, KA Lamont, WJ Tranquilli, SA Greene, SA Robertson, eds.). Ames: Wiley Blackwell.

### Critical Temperature

The critical temperature is the temperature above which a substance cannot be liquefied no matter how much pressure is applied. The critical temperature of nitrous oxide is 36.5°C. Consequently, nitrous oxide can be (and is) a liquid below this temperature, but is a gas at greater temperatures. Placing a nitrous oxide tank near a heat source will result in volatilization of liquid nitrous oxide, resulting in a high tank pressure and danger of explosion or tank venting.

The critical temperature of oxygen is −119°C, therefore at room temperature oxygen cannot be liquefied. All compressed cylinders of medical oxygen contain only gas. There are liquid oxygen tanks, but the internal tank temperature is below −119°C.

### Henry's Law

Henry's law states the solubility of a gas in a liquid is proportional to the pressure of the gas over the solution. It describes the solubility of an anesthetic in body fluids or other liquids. From it you can derive the formula  $c = k \cdot P$ , where  $c$  is the molar concentration (mol/L) of the dissolved gas and  $P$  is the pressure (in atmospheres) of the gas over the solution. For a given gas  $k$  is the Henry's law constant and is dependent on temperature.

### Partition Coefficient

A partition coefficient is the ratio of the concentration of a substance in one medium relative to another at equilibrium. It is related to the solubility of an agent. At equilibrium the partial pressure is the same throughout the body, including the alveolar gas, but the concentration of total drug may be very different due to partitioning into tissues or body fluids (Table 1.2) (Steffey et al. 2015). Partition coefficients are not absolute constants for an anesthetic

agent. Tissue composition may change as a function of age, sex, body condition, etc. and these changes may influence partitioning.

### Mechanism of Action of Inhaled Anesthetics

The simplest definition of general anesthesia is a reversible drug-induced loss of consciousness (Franks 2006). All clinically used injectable and inhaled general anesthetics reversibly induce unconsciousness at low concentrations and induce an increasing lack of responsiveness at higher concentrations (Franks 2006). Though there are a number of other desirable clinical end-points including amnesia, analgesia, immobility and muscle relaxation (Brown et al. 2010), commonly used anesthetics vary in their ability to produce these effects. Consequently, any explanation of anesthetic mechanism of action must focus on unconsciousness as the critical end-point.

Despite their longstanding and widespread use, and well-characterized pharmacodynamic effects, the specific molecular and neurophysiologic mechanisms of action of general anesthetics are only now beginning to be elucidated thanks to research accumulated over the past 120 years.

The first widely accepted theory of general anesthesia was published by Meyer and Overton in 1901. They observed that most anesthetics were lipophilic and highly hydrophobic and noted a correlation between an anesthetic's potency and the agent's oil–gas partition coefficient (Meyer 1901; Overton 1901). The so-called Meyer–Overton correlation formed the basis for numerous lipid bilayer hypotheses of anesthetic action that dominated the literature until the 1970s.

Franks and Leib were among the first to question the unitary hypothesis that the lipid bilayer was the sole site of anesthetic action (Franks & Leib 1978). They subsequently published their seminal work demonstrating that the Meyer–Overton correlation was preserved for inhibition of firefly luciferase, a soluble lipid-free model protein (Franks & Lieb 1984), which gave rise to the protein receptor hypothesis of general anesthesia. While many potential protein targets have been studied, strong evidence for direct involvement in anesthetic-induced loss of consciousness currently exists for only three: gamma-aminobutyric acid subtype A (GABA<sub>A</sub>) receptors, two-pore-domain potassium (2PK) channels, and N-methyl-D-aspartate (NMDA) receptors (Uhrig et al. 2014; Chau 2010; Franks 2008).

The GABA<sub>A</sub> family of receptors is responsible for most of the fast neuronal inhibition in the mammalian CNS and enhanced GABA-activated chloride currents cause hyperpolarization of neuronal membranes and reduce neuronal activity. Almost all general anesthetics have been found to potentiate GABA-induced chloride currents but are only

**Table 1.2** Selected partition coefficients of commonly used anesthetic agents.

| Anesthetic     | Blood:Gas Partition Coefficient | Brain:Blood Partition Coefficient |
|----------------|---------------------------------|-----------------------------------|
| Nitrous oxide  | 0.47                            | 1.1                               |
| Desflurane     | 0.42                            | 1.3                               |
| Enflurane      | 2.00                            | 1.4                               |
| Sevoflurane    | 0.68                            | 1.7                               |
| Methoxyflurane | 15.0                            | 2.0                               |
| Isoflurane     | 1.46                            | 2.7                               |
| Halothane      | 2.54                            | 2.9                               |

Source: Adapted from Steffey EP, Mama KM, Brosnan RJ. 2015. Inhalation anesthetics. In: *Veterinary Anesthesia and Analgesia: The Fifth Edition of Lumb and Jones* (KA Grimm, KA Lamont, WJ Tranquilli, SA Greene, SA Robertson, eds.). Ames: Wiley Blackwell.

capable of directly activating GABA<sub>A</sub> receptors at much higher, clinically irrelevant concentrations. Several different mutations in the alpha subunits of the GABA<sub>A</sub> receptor have been shown to eliminate or reduce the effects of volatile anesthetics (Mihic et al. 1997; Krasowski & Harrison 2000; Jenkins et al. 2001, 2002; Nishikawa & Harrison 2003; Forman & Miller 2016), whereas those in the beta subunits can reduce the effects of both intravenous and volatile anesthetic agents (Krasowski & Harrison 2000). Experiments demonstrating a correlation between stereoselectivities seen in genetically modified animals in vivo and those found with extrasynaptic GABA<sub>A</sub> receptors in vitro provide convincing evidence that these receptors play an important role in volatile anesthetic-induced loss of consciousness (Franks 2008).

The 2PK channels are thought to provide “background” modulation of neuronal excitability. Five members of this channel family (TREK1, TREK2, TASK1, TASK3, and TRESK) can be directly activated by volatile anesthetics (Patel et al. 1999) but anesthetic sensitivity among these channels is not uniform. Anesthetic activation of 2PK channels generally inhibits neuronal activity by either hyperpolarizing the membrane and/or increasing membrane conductance, thus reducing the effects of excitatory currents (Franks 2008). However, these channels are also found presynaptically and their activation here may be inhibitory (if at an excitatory synapse) or excitatory (if at an inhibitory synapse). A growing body of evidence suggests these channels mediate at least some of the effects of volatile anesthetics (Franks 2008).

NMDA receptors mediate the slow components of synaptic transmission in the CNS and may be important targets for certain anesthetics. Both gaseous (i.e., xenon and nitrous oxide) and volatile (i.e., isoflurane, sevoflurane, halothane, and desflurane) inhaled anesthetics inhibit NMDA receptors to some extent but considerable variability exists among agents. Although it is likely that NMDA receptor antagonism plays a role in the actions of xenon and nitrous oxide and, to a lesser extent, the volatile agents, additional targets are required to account for their ability to cause loss of consciousness (Franks 2008).

Determining the cascade of events that follow binding of an anesthetic agent to its molecular target(s) leading to loss of consciousness has proven difficult. Remarkable advances in neuroscience and neuroimaging techniques (e.g., positron emission tomography, functional magnetic resonance imaging [fMRI], magnetoencephalography, and electroencephalography) over the past two decades are beginning to unravel the neural correlates of consciousness which will help answer this longstanding question.

Convergent evidence from multiple studies involving multiple species, multiple neuroimaging modalities,

multiple analytic techniques, and diverse drug classes suggests that anesthetic-induced unconsciousness is characterized by a functional fragmentation (also referred to variably as a disconnection or uncoupling) of cortical and thalamocortical networks (Hudetz & Mashour 2016; Li et al. 2017; Mashour & Hudetz 2017). One specific cortical region, known as the lateral frontoparietal network, has received considerable attention over the last decade. Both fMRI and electroencephalographic studies of diverse classes of anesthetics, including volatile inhaled agents, consistently show a functional breakdown in frontoparietal connectivity and surrogates of frontoparietal information transfer (Hudetz & Mashour 2016; Mashour & Hudetz 2017). As our understanding of consciousness continues to evolve, traditional references to the “depth of anesthetic hypnosis” may require revision to reflect the fact that general anesthetics may variably suppress certain higher-order components of consciousness while retaining others (Sleigh et al. 2018).

## **Inhalant Anesthetic Pharmacokinetics**

### ***Anesthetic Uptake and Distribution***

A series of partial pressure gradients, beginning at the vaporizer, continuing in the anesthetic breathing circuit, the airways, alveoli, blood, and ending in the tissues will drive the movement of an anesthetic gas. The movement of that gas will continue until equal partial pressures are present throughout the system. Since the lung is the point of entry and exit to the body, the alveolar partial pressure governs the partial pressure of the anesthetic in all body tissues. Therefore, it is most important to understand how to influence the alveolar partial pressure. Increasing alveolar minute ventilation, flow rates at the level of the vaporizer, and inspired anesthetic concentration can speed the delivery of anesthetic and increase the rate of rise of alveolar anesthetic partial pressure. Solubility, cardiac output, and the alveolar-to-venous anesthetic gradient are factors that determine the uptake of the anesthetic from the alveoli into the blood. Solubility describes the affinity of the gas for a medium such as blood or adipose tissue and is reported as a partition coefficient. The blood/gas partition coefficient describes how the gas will partition itself between the two phases (blood and alveolar gas) after equilibrium has been reached. Isoflurane, for example, has a blood/gas partition coefficient of approximately 1.46 (Steffey et al. 2015). This means that if the gas partial pressures are in equilibrium, the concentration in blood will be 1.46 times greater than the concentration in the alveoli. A higher blood/gas partition coefficient means a greater uptake of the gas into the blood, therefore a slower rate of rise of alveolar and blood partial pressure. Since the blood partial pressure rise is slower, it takes longer for the brain partial pressure of the gas to increase, resulting in a longer induction time.

Increased cardiac output exposes the alveoli to more blood per unit time. The greater volume of blood removes more inhalant anesthetic from the alveoli, therefore lowering the alveolar partial pressure. The agent might be distributed faster within the body, but the partial pressure in the arterial blood is lower. It will take longer for the gas to reach equilibrium between the alveoli and the brain, therefore a high cardiac output usually prolongs induction time. The alveolar to venous partial pressure difference reflects the tissue uptake of the inhaled anesthetic. A large difference is caused by increased uptake of the gas by the tissues during the induction phase.

Transfer of the gas from the arterial blood into tissues such as the brain will depend on perfusion and the relative solubility of the gas in the different tissues. The brain/blood coefficient describes how the gas will partition itself between the two phases after equilibrium has been reached. Isoflurane has a brain/blood coefficient of 2.7, therefore when the system is at equilibrium the concentration in the brain will be 2.7 times greater than the concentration in the blood (Steffey et al. 2015). All contemporary inhalation anesthetics have high adipose/blood partition coefficients. This means that most of the gas will accumulate in adipose tissue as time goes by. The partial pressure of the gas in adipose tissue will rise very slowly since this tissue has a high capacity (as indicated by the high adipose/blood partition coefficient). Inhalation anesthetics stored in obese patients may delay awakening at the end of long periods of anesthesia. Fortunately, adipose tissue has a relatively low blood flow and doesn't accumulate significant amounts of anesthetic during the short periods of anesthesia commonly encountered in veterinary medicine.

### ***Elimination of Inhaled Anesthetics***

The rate of induction and recovery from anesthesia with inhalant anesthetics differs between agents due to differences in tissue solubility, but general statements can be made. During induction all tissue partial pressures are zero. During recovery, different tissues in the body have different partial pressures of anesthetic, which are governed by the tissue anesthetic content and not the alveolar partial pressure. Recovery is not as controllable as induction of anesthesia. During recovery from anesthesia, elimination occurs due to exhalation and biotransformation.

The enzymes responsible for inhalant anesthetic metabolism are mainly located in liver and kidneys. Anesthetic elimination via metabolism is approximately 50% for methoxyflurane, 10–20% for halothane, 5–8% for sevoflurane, 2.5% for enflurane, about 0.2% for isoflurane, 0.001% for desflurane, and nearly zero for nitrous oxide (Steffey et al. 2015). The amount of anesthetic eliminated from the body during anesthesia due to metabolism is small

compared with the amount exhaled. However, anesthetic metabolism accounts for a larger proportion of the anesthetic clearance after anesthetic delivery ceases. The low, but prolonged, blood partial pressure of the anesthetic found after terminating delivery is no longer overwhelming the enzyme systems (enzymes become saturated above ~1 MAC [minimum alveolar concentration]) so metabolism accounts for a larger proportion of clearance than it did during exposure to high partial pressures.

Elimination of the anesthetic via the lungs can be complex. The first point to consider is what effect an increase in alveolar minute ventilation will have on recovery. During recovery, increasing minute ventilation will decrease alveolar anesthetic partial pressure and increase the gradient for diffusion from the blood to the alveoli. This increases elimination, especially for most anesthetic agents with high blood/gas partition coefficients.

Another situation to consider is what effect a change in cardiac output will have on the rate of decrease of partial pressure of the inhalant anesthetic. During induction, high cardiac output will increase the rate at which anesthetic is removed from the lung, slowing the rate of rise of anesthetic partial pressure and slowing induction. When cardiac output is reduced (e.g., cardiogenic shock) there is a slower removal of anesthetic and subsequently a faster rate of rise of alveolar partial pressure and induction occurs. During recovery a high cardiac output will increase the rate at which anesthetic is returned to the lung for excretion. Since the partial pressure of anesthetic in the blood is determined by the tissues, the higher blood flow will shorten recovery. During low cardiac output situations there will be a slower recovery due to the decreased rate at which tissue anesthetic partial pressure decreases.

The last major influence on the rate of induction and recovery is the solubility of the anesthetic agent. Agents with high blood/gas solubility will be partitioned into the blood to a greater extent than agents with low blood/gas solubility. The blood acts as a depot for agent maintaining anesthetic partial pressure. Agents with low blood/gas solubility do not partition into the blood to the extent of highly soluble agents, thus the decrease in partial pressure is faster and recovery time is reduced. Highly soluble agents have high blood concentrations and it will take longer for the partial pressure to decrease if all other factors are equal. In summary, elimination of a volatile anesthetic depends on the ventilation, cardiac output, and solubility of the gas in blood and tissue.

### **Control of the Partial Pressure of Delivered Anesthetic**

Inhalant anesthetics can be classified as either gaseous (nitrous oxide and xenon) or volatile (isoflurane, sevoflurane, halothane, methoxyflurane, and desflurane). Gaseous

anesthetics are usually delivered to the anesthesia machine under pressure and their rate of delivery to the breathing circuit is controlled by a flow meter. Volatile anesthetics are liquids at room temperature and pressure, and are usually delivered by a specialized apparatus that controls the volatilization of the liquid and proportioning of the vapor in the fresh gas delivered to the patient. A vaporizer can be as simple as piece of cotton soaked with agent held near the nose (not recommended) or can be as complex as the desflurane vaporizer, which is an electrically powered device with a heated vaporizing chamber.

### The Breathing System

With most modern anesthetic machines the outflow gas from the vaporizer will be delivered to the patient through a set of tubes and machinery collectively called a breathing system. There are many styles of breathing systems, each with a multitude of uses. It is important that the anesthetist understands how the type of breathing circuit used will impact the rate at which the anesthetic concentration can be changed and the relationship between the vaporizer setting and inspired concentration.

### Waste Anesthetic Gases

The health effects of chronic exposure to waste anesthetic gases are not completely known. The frequency of inhalant anesthetic use and the lack of significant associations between exposure and most types of chronic toxicities (e.g., cancer, infertility, birth defects, etc.) would suggest there is only a very low risk (if any) associated with chronic exposure. However, certain individuals are highly susceptible to potentially life-threatening reactions, even with trace level exposure (e.g., malignant hyperthermia). In light of this, and with the admission that we do not completely understand all the risks associated with chronic exposure, it is generally agreed that the exposure of personnel be kept as low as reasonably acceptable (ALARA). In the United States, the Occupational Safety and Health Administration requires veterinary hospitals to maintain a system to prevent waste gases from building up in the area of use and can enforce exposure limits that are consistent with guidance from the National Institute of Occupational Safety and Health (NIOSH). Currently, no NIOSH-recommended exposure limits exist for the three most currently used anesthetics (isoflurane, desflurane, and sevoflurane) and at present NIOSH has no published permissible exposure limits regulating these specific agents (American Veterinary Medical Association 2020).

### Minimum Alveolar Concentration

Although loss of consciousness is arguably the most critical end-point of general anesthesia (see section on the

inhaled anesthetic mechanism of action), it remains challenging to measure in a clinical setting. Immobility in the face of a noxious stimulus is also an important end-point and has formed the basis of the standard unit of inhaled anesthetic potency – minimum alveolar concentration (MAC) – since the 1960s (Merkel & Eger 1963). While our understanding of how inhaled anesthetics produce immobility continues to evolve (Eger et al. 2008), MAC remains the standard measurement of dose in clinical anesthesia. It is defined as the minimum alveolar concentration at 1 atmosphere required to prevent gross purposeful movement in 50% of the subjects tested, following a 60-second application of a supramaximal stimulus (Steffey et al. 2015). One MAC is by definition the  $EC_{50}$  (i.e., the effective concentration in 50% of patients) for that agent. Animals awaken from anesthesia at approximately 0.5 MAC, surgical anesthesia occurs at approximately 1.3 MAC, and severe autonomic nervous system depression occurs around 2 MAC.

Birds and many reptiles do not have true alveoli so the concept of MAC has been modified or redefined to be the minimum anesthetic concentration. It is not identical to MAC from other species, but closely approximates it in many ways.

### Physiological and Pharmacological Factors That Alter MAC

Minimum alveolar concentration is age-dependent, being lowest in newborns, reaching a peak in infants, and then decreasing progressively with increasing age (Lerman et al. 1983, 1994; Taylor & Lerman 1991). Increases in MAC can also occur from hyperthermia and hypernatremia, and decreases in MAC can result from hypothermia, hyponatremia, pregnancy, hypotension, and drugs such as lithium, lidocaine, opioids, and alpha-2 adrenergic agonists.

### General Pharmacological Actions of Inhalant Anesthetics

Inhalant anesthetic agents have more similarities than differences with respect to their effects on vital organ systems. The differences are primarily related to the speed and magnitude with which the changes occur. There are a few classic differences that have been included in the following synopsis.

#### Central Nervous System

All inhalant general anesthetics alter consciousness, memory, and pain perception by acting on the CNS. Most inhalant anesthetics cause a mild to moderate decrease in the cerebral metabolic requirement for oxygen ( $CMRO_2$ ) and they usually have minimal effects on cerebral blood flow autoregulation at low MAC multiples (Mielck et al.

1998, 1999). Patients with intracranial hypertension should not be anesthetized with nitrous oxide because it may cause an increase in CMRO<sub>2</sub> (Roald et al. 1991; Algotsson et al. 1992; Hoffman et al. 1995). Halothane is also a poor choice because of its significant effects on cerebral blood flow autoregulation (Steffey et al. 2015). Isoflurane, sevoflurane, and desflurane are the inhalants of choice.

### Cardiovascular System

Most inhalant anesthetic agents cause direct myocardial depression. Halothane is the most depressant on contractility, but it generally has the fewest effects on vascular resistance (Steffey et al. 2015). Isoflurane, enflurane, sevoflurane, and desflurane cause some degree of vasodilatation, which tends to improve forward blood flow and maintain tissue perfusion. The reduction in afterload also tends to offset some of the direct myocardial depressant effects and may result in a net improvement in cardiac output. Nitrous oxide is a sympathomimetic and can improve contractility, blood pressure, and heart rate at light levels of anesthesia. Rapid changes in anesthetic concentration (especially with desflurane) may result in a sympathetic response and temporarily increase cardiac work.

### Respiratory System

All anesthetics tend to depress the chemoreceptor response to carbon dioxide, leading to an accumulation of carbon dioxide and a respiratory acidosis unless ventilation is assisted or controlled. The ether derivatives tend to be the most depressant, but all agents may cause significant depression. Most inhalant agents may interfere with hypoxic pulmonary vasoconstriction and may worsen ventilation-perfusion matching in the lung. This is most dramatic in larger animals, where significant pulmonary shunting is often observed.

### Genital-Renal Systems

Most anesthetics cause a decrease in renal perfusion and an increase in antidiuretic hormone (ADH) secretion. Inhalant anesthetics may be the safest anesthetic techniques in anuric renal failure since pulmonary excretion is not dependent on renal function.

Inhalant anesthetics may cause an increase in postpartum uterine bleeding. This is a bigger consideration in primate anesthesia due to placentation characteristics. Isoflurane, sevoflurane, desflurane, and nitrous oxide have been advocated for use during Caesarian section because of the rapid onset and termination of effect, and the transient effects on the delivered fetuses. Methoxyflurane and halothane are less desirable due to their greater solubility and slower elimination.

## Clinically Useful Inhalant Anesthetics

### Nitrous Oxide

Nitrous oxide is commonly used in combination with a primary inhalant or injectable anesthetic drug. The reason it is not useful in veterinary anesthesia as a solo anesthetic is because of its low potency. Nitrous oxide's MAC value has been estimated to be near 100% for humans and closer to 200% for veterinary patients. It is obvious that 200% nitrous oxide cannot be delivered; in fact no more than 79% nitrous oxide can be safely delivered without creating a hypoxic gas mixture. In practice it is common to use a 50% nitrous oxide mixture with the balance of the mix being oxygen. If 50% nitrous oxide is delivered to an animal it is only providing approximately 0.25 MAC of anesthesia. A potent volatile anesthetic, injectable agent, or other sedative/analgesic drug must supply the remaining 0.75 MAC. Because of this limited anesthetic effect, nitrous oxide use for anesthetic maintenance is not widespread in veterinary medicine. Nitrous oxide is used by some anesthetists during induction of anesthesia for the *second gas effect*. Since nitrous oxide is present in the inspired gas mixture in a relatively high concentration and it rapidly diffuses into the body from the alveoli, the rate of rise of partial pressure of a second coadministered inhalant anesthetic is increased and the induction time can be shortened.

Nitrous oxide has a low blood/gas partition coefficient and has a rapid onset and recovery. The gas can diffuse out of the blood so rapidly that if nitrous oxide delivery is suddenly halted and supplemental oxygen is not administered, a situation known as *diffusion hypoxia* may result. Diffusion hypoxia happens when the mass movement of nitrous oxide down its partial pressure gradient results in high alveolar nitrous oxide partial pressure at the expense of oxygen and nitrogen partial pressures. Since breathing room air will result in an alveolar oxygen partial pressure of approximately 100 mmHg under ideal circumstances, any displacement of oxygen by nitrous oxide will result in alveolar hypoxia. Diffusion hypoxia can be minimized or prevented by continuing the administration of oxygen enriched gas for 5–10 minutes following the discontinuation of nitrous oxide. This helps because during normal breathing, 100% oxygen should result in an alveolar oxygen partial pressure close to 500 mmHg. The partial pressure of oxygen can drop a lot further before hypoxia develops.

Nitrous oxide is contraindicated in animals with pneumothorax, gastric dilatation/rumen tympany, gas embolism, and other conditions that are exacerbated by accumulation of gas inside a closed space. This effect is caused by diffusion of nitrous oxide out of the blood into the preexisting gas space in an attempt to establish equilibrium. Nitrous oxide is also contraindicated in animals with

gas diffusion impairment such as interstitial pneumonia. These animals typically have low arterial oxygen partial pressure when breathing oxygen-rich mixtures. The dilution of oxygen by nitrous oxide will lower the inspired oxygen partial pressure and may worsen hypoxemia.

### Halothane

Halothane was a major advancement in inhalant anesthesia in its day. It was introduced in the late 1950s and was potent, nonirritating, and nonflammable. Chemically it is classified as a halogenated hydrocarbon and it is not chemically related to the ethers. Halothane was used widely in human anesthesia until it became apparent there were potentially fatal adverse effects associated with its use. Human patients developed a syndrome known as *halothane hepatitis* (Daghfous et al. 2003; Neuberger 1998). This rare, but life-threatening, complication is still somewhat of a mystery, although an immunological etiology is implicated. The disease appears as a fulminant hepatitis, similar to that seen with viral hepatitis, which develops after a short period of apparent recovery. A second, more common, form of hepatitis is less severe and is characterized by a reversible elevation in liver enzymes. The etiology of this second form is thought to be anesthetic-related hepatic hypoxia and does not appear to be immune related. Diagnosis of the correct form is important since repeated exposure to halothane, or any of the volatile agents producing trifluoroacetic acid, is more likely to trigger the immunologically mediated form and result in high morbidity and mortality. Both forms are not commonly documented in veterinary patients; however, transient elevation of liver enzymes may occur postoperatively in some patients. A thorough diagnostic workup is required due to the non-specific and multifactorial etiology of elevated liver enzymes.

A second complication associated with halothane anesthesia is the development of arrhythmias. Halogenated hydrocarbon anesthetics, especially halothane, can sensitize the myocardium to the arrhythmogenic effects of epinephrine. Halothane is generally contraindicated in patients that are predisposed to ventricular arrhythmias (e.g., hypoxia, trauma, or myocardial disease) (Steffey et al. 2015). Arrhythmias that develop during halothane anesthesia may resolve when the anesthetic agent is switched to isoflurane or sevoflurane. Other causes of peri-anesthetic arrhythmias should also be ruled out.

Halothane undergoes extensive hepatic metabolism (~20%) and is not chemically stable (Steffey et al. 2015). Commercially available halothane contains thymol, a preservative, which does not volatilize to the same degree as

halothane. This results in a sticky residue inside the vaporizer that should be cleaned out during periodic maintenance. Veterinary use of halothane is declining due to the increasing popularity of isoflurane and sevoflurane, and its limited availability worldwide.

### Isoflurane

Isoflurane is arguably the most widely used veterinary inhalant anesthetic in the world today. It is stable, potent, and undergoes little metabolism. Isoflurane can be irritating to airway tissues at high-inspired concentrations and its use for induction in people has been limited because of patient complaints and complications. However, in veterinary medicine isoflurane mask induction is still common. Isoflurane is a potent agent (MAC ~1.3% in dogs) and has a high saturated vapor pressure (~240 mmHg at room temperature) (Steffey et al. 2015). These characteristics, coupled with the fact that it is possible to cause rapid partial pressure changes in the brain, would suggest that only precision vaporizers located outside the circuit should be used to deliver the agent. However, several reports of the use of modified vaporizer inside circuit (VIC) vaporizers suggest that this type of anesthetic system can be used to safely administer the agent (Bednarski et al. 1993; Laredo et al. 1998).

Isoflurane metabolism is minimal (less than 1%) and fluoride-induced nephrotoxicity is uncommon. Isoflurane and many of the ether-derivative volatile agents are excellent vasodilators and can cause or worsen hypotension.

### Desflurane

Desflurane use in veterinary medicine is limited to academic institutions and a very limited number of private practices. The main disadvantage to desflurane use is the cost associated with the agent and the cost associated with a specialized vaporizer that is required to deliver the drug. Desflurane is extremely insoluble and is capable of producing extremely rapid inductions and recoveries (Barter et al. 2004; Clarke 1999). Its main market is for human outpatient anesthesia, where rapid recovery is a large cost savings. It is highly fluorinated, has a very low potency (MAC ~9%), and has a high saturated vapor pressure (~670 mmHg at room temperature) (Steffey et al. 2015). Desflurane boils at 23°C and must be handled using specialized apparatus for vaporizer filling. The vaporizer is specific for desflurane and is electrically heated to boil the desflurane so that a reliable vapor pressure will be produced. Then sophisticated differential pressure transducers and electronic circuits calculate an injection ratio for delivery of the desired anesthetic concentration. Desflurane is very stable and undergoes almost no metabolism.

### Sevoflurane

Sevoflurane is the newest volatile inhalant anesthetic approved for veterinary use. Sevoflurane has a low blood/gas partition coefficient ( $\sim 0.7$ ) which is greater than desflurane and nitrous oxide, but about half of that of isoflurane. Extensive pulmonary elimination of sevoflurane minimizes the amount available for metabolism. Up to 3–8% of the sevoflurane dose is metabolized and appears in the urine as inorganic fluoride (Steffey et al. 2015). This fluoride exposure does not appear to be clinically significant, although serum levels of fluoride can approach those previously reported to be nephrotoxic for methoxyflurane. Factors other than peak serum fluoride concentrations appear important for predicting the incidence of nephrotoxicity (Driessen et al. 2002).

Sevoflurane represents a deviation from the methyl ethyl ether structural theme present in other contemporary volatile anesthetics. Sevoflurane is chemically related to methyl-isopropyl ethers. The structure is significant because an important metabolite of most methyl-ethyl ether volatile anesthetic agents (trifluoroacetic acid) is a suspected trigger of halothane hepatitis. Sevoflurane cannot be metabolized to form this compound. This is not a major consideration in veterinary medicine but is important in human anesthesia. Sevoflurane is also pleasant and nonirritating when used for mask induction and many pediatric anesthesiologists suggest this agent is the drug of choice for pediatric induction via mask. Sevoflurane is less potent than isoflurane (MAC  $\sim 2.3\%$  for dogs and horses). When used for induction of anesthesia it is common to use 7–9% sevoflurane.

An early subject of controversy surrounding sevoflurane anesthesia was the production of compound A. Compound A is a degradation product produced when sevoflurane reacts with the carbon dioxide absorbent. Early toxicology studies performed in rats suggested that proximal tubular renal damage could result from clinically relevant exposure to compound A. This led to the suggestion that sevoflurane should not be used in closed-circuit anesthesia or with fresh gas flow rates lower than 2 L per minute. However, since that time little clinical evidence of renal damage in humans and dogs has emerged, even with very low fresh gas flows. Some have suggested that rats have a 10–100 times higher level of the enzyme beta-lyase that is believed to convert the intermediate compounds of Compound A metabolism to a nephrotoxic molecule (Kharasch et al. 2005; Sheffels et al. 2004). Humans and dogs do not appear to have the same level of enzyme conversion and are therefore less susceptible to Compound A toxicity. Safety studies in most other rodents and exotic animals are not published and caution should be used when

**Table 1.3** Structure and characteristics of inhalation anesthetics.

| Agent          | Year Introduced | Structure                                 | Type   |
|----------------|-----------------|-------------------------------------------|--------|
| Halothane      | 1956            | $\text{CF}_3\text{-CHClBr}$               | Alkane |
| Isoflurane     | 1981            | $\text{CF}_3\text{CHCl-O-CHF}_2$          | Ether  |
| Enflurane      | 1972            | $\text{CHClF-CF}_2\text{-O-CHF}_2$        | Ether  |
| Methoxyflurane | 1960            | $\text{CHCl}_2\text{-CF}_2\text{-O-CH}_3$ | Ether  |
| Desflurane     | 1992            | $\text{CF}_3\text{CHF-O-CHF}_2$           | Ether  |

administering sevoflurane via a breathing system using a carbon dioxide absorbant until further safety data are available (Table 1.3).

## Injectable Anesthetics

Injectable anesthetics are an important family of compounds used for immobilization and anesthesia of wildlife. The dissociative anesthetics in particular are commonly combined with other adjunctive drugs such as alpha-2 adrenergic agonists and opioids to improve the reliability and speed of onset of action.

### Barbiturates

Barbiturates can be classified in several ways. One is by chemical structure. Oxybarbiturates are historically important, but not commonly used today due to their slower onset of action, long recovery characteristics, and relatively small margin of safety. Pentobarbital is the prototypical oxybarbiturate. It has been combined with several adjunctive drugs for anesthesia. The thio (i.e., sulfur-substituted) analog of pentobarbital, thiopental, was commonly used in veterinary medicine for intravenous induction of anesthesia for many years but is no longer widely available.

Barbiturates cause anesthesia through global depression of CNS activity. This is accomplished through interference with nervous system impulse conduction. Like many other anesthetics, other excitable tissues can be affected, resulting in commonly encountered side effects, including depression of cardiorespiratory function. Barbiturates decrease cerebral blood flow (CBF) and the  $\text{CMRO}_2$ .  $\text{CMRO}_2$  decreases progressively until electroencephalographic activity becomes isoelectric (Branson 2007).

## Propofol

Propofol (2,6-di-isopropylphenol) is commonly used for sedation, induction, and maintenance of anesthesia in humans and domestic species. Propofol is supplied as a milky white liquid for intravenous injection. It is insoluble in aqueous solution therefore it is usually formulated as an emulsion of 10% soybean oil, 2.25% glycerol, and 1.2% egg phosphatide. Some formulations of propofol (e.g., Diprivan®, Propoflo™) do not contain preservative and will support bacterial and fungal growth should the drug become contaminated. This has led to the label recommendation of discarding unused drug at the end of the procedure or within 6 hours of opening a vial. Other formulations have additives such as benzyl alcohol (e.g., Propoflo28®) to improve stability or reduce the potential of contamination with storage. Species-sensitivities to these additives should be investigated before their use (Davidson 2001).

Propofol is classified as an ultrashort-acting injectable anesthetic agent. Duration of effect is typically 5–10 minutes in dogs and 5–20 minutes in cats. Its rapid recovery characteristics are maintained in most species following prolonged infusions. Recovery times may be prolonged in the cat (and other species that have reduced capacity for glucuronidation of drugs) following repeated doses or continuous rate infusions.

Propofol has been used in dogs, cats, horses, pigs, goats, sheep, and even birds. Wild turkeys, mallard ducks, pigeons, and chickens have been anesthetized with propofol, but there is significant cardiorespiratory depression in ducks and chickens, indicating birds may need ventilatory support during anesthesia (Machin & Caulkett 1998). Apnea and respiratory depression are the best known side effects of propofol administration. The incidence of apnea may be reduced by administering the drug over 60–90 seconds (Muir & Gadawski 1998). It would be prudent to be prepared to intubate and support ventilation if apnea occurs. Pain is reported on propofol injection by some people. Muscle fasciculations and spontaneous twitching can occur in some animals.

## Dissociative Anesthetics

### Ketamine

Most veterinary formulations of ketamine are a racemic mixture consisting of two optical enantiomers. However, in many countries (*S*)-ketamine is available as a human or veterinary product. The *S* enantiomer is less cardiodepressant and has a four-fold greater affinity for the phencyclidine site in the NMDA receptor. Serotonin transport is inhibited two-fold by the *R* form. Some of ketamine's effects are not stereoselective. Norepinephrine release is equivalent from the *S* and *R* forms (Kohrs & Durieux 1998).

Ketamine can be administered intramuscularly to anesthetize animals which are not easily given drugs intravenously. Intramuscular administration will produce a longer duration of anesthesia than intravenous administration, but the recovery is usually longer and can be accompanied by more dysphoria. Recovery from ketamine appears to be due to redistribution and metabolism similar to the thiobarbiturates. Hepatic biotransformation to norketamine (metabolite I) and dehydronorketamine (metabolite II) is the major route of metabolism in most species studied. It was thought ketamine was excreted unchanged in the urine of cats, but this originated from one paper published in 1978 by Gaskell et al. and since that time it has been shown by Waterman that biotransformation is an important route of elimination in domestic cats (Waterman 1983). Norketamine is about one-third to one-fifth as potent as the parent compound but may contribute to the prolonged analgesic effects of ketamine (Kohrs & Durieux 1998).

Ketamine produces a form of anesthesia that is different from other hypnotic drugs. In general terms, ketamine induces anesthesia and amnesia by functional disruption (dissociation) of the CNS through marked CNS stimulation resulting in catalepsy, immobility, amnesia, and marked analgesia. Electroencephalographic analysis indicates that depression of the thalamocortical system occurs in conjunction with activation of the limbic system. Awakening from ketamine anesthesia in people is frequently characterized by disagreeable dreams and hallucinations. Sometimes these unpleasant occurrences may recur days or weeks later. Almost half of adults over the age of 30 exhibit delirium or excitement, or experience visual disturbances. The occurrence of adverse psychological experiences is much lower in children. The incidence of adverse psychological experiences in animals is unknown, but a significant number of animals transiently vocalize and have motor disturbances during recovery.

Ketamine's neuropharmacology is complex. The compound interacts with NMDA and non-NMDA glutamate receptors, nicotinic, muscarinic cholinergic, monoaminergic, and opioid receptors. In addition there are interactions with voltage-dependent ion channels such as Na<sup>+</sup> and L-type Ca<sup>2+</sup> channels. It is believed that the NMDA receptor antagonism accounts for most of the analgesic, amnesic, psychomimetic, and neuroprotective effects of the compound, but the exact mechanism of its anesthetic action is not known. NMDA receptor activation is believed to play a role in the "memory" of the CNS, which is involved in the "wind-up," hyperalgesia, and allodynia seen in certain pain syndromes (Kohrs & Durieux 1998).

Ketamine can increase the CMRO<sub>2</sub> due to increased metabolic activity associated with increased activity in certain

areas of the brain. Intracranial pressure (ICP) also increases, possibly because of two mechanisms: (1) ketamine can increase mean arterial blood pressure so CBF can increase and ICP can passively increase in patients with altered autoregulation, and (2) ketamine can depress respiration, increasing the arterial partial pressure of carbon dioxide  $\text{PaCO}_2$ . The brain responds to elevations in  $\text{PaCO}_2$  by increasing CBF, which will increase ICP. Ventilation may reduce the increase in CBF. Current clinical dogma dictates avoiding ketamine in patients with suspected head trauma.

Ketamine causes a characteristic breathing pattern termed *apneustic breathing*, characterized by prolonged inspiratory duration and relatively short expiratory time. When ketamine is administered by itself, it typically causes minimal respiratory depression that is short-lived. Hypoxic and hypercapnic respiratory regulation appears to remain intact, but ketamine is seldom given alone. It is often combined with benzodiazepines, acepromazine, opioids, or alpha-2 adrenergic agonists. The combined effect of these drugs is usually decreased minute ventilation, increased  $\text{PaCO}_2$ , and mild respiratory acidosis.

Ketamine, when given to animals with functioning sympathetic nervous systems, generally increases heart rate and arterial blood pressure. Cardiac output will usually stay the same or slightly increase. Ketamine is seldom given alone to healthy animals. The use of adjunctive drugs, such as benzodiazepines, acepromazine, or alpha-2 adrenergic agonists, tends to blunt the sympathomimetic effect of ketamine and will tend to decrease cardiac function and decrease arterial blood pressure.

### Tiletamine/Zolazepam

Tiletamine/zolazepam combinations are available in a fixed ratio. Telazol® is a non-narcotic, nonbarbiturate, injectable anesthetic agent. Chemically, Telazol® is a combination of equal parts by weight of tiletamine hydrochloride (2-[ethylamino]-2-[2-thienyl]-cyclohexanone hydrochloride), an arylaminocycloalkanone dissociative anesthetic, and zolazepam hydrochloride (4-[o-fluorophenyl]-6,8-dihydro-1,3,8-trimethylpyrazolo[3,4-e][1,4]diazepin-7[1H]-1-hydrochloride), a benzodiazepine having minor tranquilizing properties. The product is supplied sterile in vials, each containing a total of 500 mg of active drug as free base equivalents and 288.5 mg of mannitol. The addition of 5 mL of diluent produces a solution containing the equivalent of 50 mg of tiletamine base, 50 mg of zolazepam base and 57.7 mg of mannitol per milliliter. The resulting solution has a pH of 2–3.5. Zoletil® is available in many countries outside North America and is commonly marketed as a mixture containing 25 mg/mL each of zolazepam and tiletamine (Zoletil 50®) or 50 mg/mL each (Zoletil 100®).

Duration of effect is dependent on the route of administration and amount of drug given. When used intravenously it lasts approximately 15–20 minutes. When given intramuscularly it may last 30–45 minutes. It is commonly used in place of ketamine and its duration is typically longer.

Tiletamine induces dissociative anesthesia similar to ketamine. It has the potential to cause seizure activity, but when combined with zolazepam the incidence of seizures is greatly reduced. Its effects on CBF and ICP are similar to those of ketamine. Nephrotoxicity in New Zealand white rabbits has been reported following Telazol® administration (Doerning et al. 1992). Anecdotally, tigers do not appear to recover well after tiletamine/zolazepam therefore its use is generally contraindicated. Tiletamine/zolazepam can be combined with other drugs to improve its analgesic and recovery characteristics.

## Miscellaneous Anesthetics

### Etomidate

Etomidate has been used extensively as a hypnotic agent for the induction of anesthesia in humans, but less commonly in other species. Etomidate is a rapidly acting, ultra-short acting imidazole derivative. The duration of effect following intravenous bolus administration is typically 5–10 minutes. Etomidate causes dose-dependent CNS depression leading to sedation, hypnosis, and finally an isoelectric electroencephalogram.

Etomidate, in contrast to almost all other induction agents, does not seem to cause significant depression of cardiac contractility and has minimal effects on heart rate, cardiac output, and arterial blood pressure. Elimination of etomidate occurs by ester-hydrolysis in plasma and in the liver at approximately equal rates. Metabolism of etomidate in the liver is a capacity-limited Michaelis–Menten process. Hepatic hydrolysis results in the corresponding inactive carboxylic acid. Etomidate will temporarily reduce steroidogenesis (Boidin 1985; Moon 1997). Steroid synthesis usually increases with the stress of anesthesia so the net effect may be little or no change (Dodam et al. 1990). It is not a clinical contraindication except for animals with hypoadrenocorticism (Addison's disease). Intravenous administration of etomidate may induce excitement, myoclonus, pain on injection, vomiting, and apnea during induction of anesthesia. Some animals may have purposeless myoclonic muscle movements during recovery from anesthesia. The frequency and severity of the side effects can be attenuated or eliminated by the administration of adjunctive drugs such as diazepam, acepromazine, or opioids prior to etomidate administration. A constant rate infusion of etomidate may result in hemolysis (Van de Wiele et al. 1995; Moon 1994). This is thought to be due to

the propylene glycol carrier and the very high osmolality of available products (Doenicke et al. 1997).

### Alfaxalone

#### *Alfaxan*®

Alfaxalone is a synthetic neurosteroid anesthetic with a relatively wide margin of safety and little cardiovascular or respiratory depression. Due to its poor water solubility, earlier formulations (*Saffan*®) were a mixture of alfaxalone and alfadolone combined with cremophor EL® as a solubilizing agent. Adverse effects, including histamine release and anaphylactic reactions associated with Cremophor, led to the withdrawal of that product from the market. A novel formulation of alfaxalone, *Alfaxan-CD*®, was released in Australia in 2001, in the United Kingdom in 2007, in Europe in 2008, in Canada in 2011, and in the United States in 2012. It is a 1% aqueous solution with a pH of 6.5–7.0 containing a non-cremophor vehicle (2-hydroxypropyl- $\beta$ -cyclodextrin) which does not cause histamine release. The drug is approved for induction and maintenance of general anesthesia in dogs and cats via intravenous administration and has also been approved for intramuscular administration in some countries. Until recently, *Alfaxan*® was exclusively supplied as a 10-mL vial without added antimicrobial preservatives. This resulted in a shelf-life of only 6 hours and a label recommendation to discard unused drug within this timeframe (Berry 2015). A newer multi-dose formulation containing the preservatives ethanol, chlorocresol, and benzethonium chloride has recently been approved and appears to have the same efficacy and safety as the preservative-free formulation but offers the advantage of a 28-day shelf-life.

Like other injectable anesthetics, alfaxalone produces its clinical effects via potentiation of GABA at GABA<sub>A</sub> receptors in the CNS, resulting in modulation of chloride transport and neuronal hyperpolarization (Berry 2015). After intravenous administration, it produces rapid onset of action and satisfactory muscle relaxation but has no significant analgesic effects. In dogs and cats, it undergoes rapid hepatic metabolism, has a relatively short half-life, and does not accumulate after repeated dosing (Ambros et al. 2008; Ferre et al. 2006; Whittam et al. 2008).

Depending on the species and country-specific labelling, alfaxalone can be used for intramuscular sedation or premedication, for intravenous induction of anesthesia prior to transfer to an inhaled agent, or for intravenous maintenance of anesthesia via intermittent boluses for short procedures or via continuous rate infusion (CRI) for longer procedures. Other off-label routes of administration have been reported, including subcutaneous injection in various domestic and exotic species, intraperitoneal injection in laboratory animals, and immersion/irrigation in fish and amphibians.

Premedication or coadministration with a sedative and/or analgesic is recommended to reduce the dose of alfaxalone required, provide analgesia, and smooth recovery, which can otherwise be rough. Paddling, vocalization, rigidity, myoclonus, and trembling have been reported in dogs and cats recovering from alfaxalone that did not receive other drugs concurrently (Berry 2015). Also, with intramuscular administration, the relatively large volume of alfaxalone required can be a limiting factor and coadministration of more potent sedative and/or analgesic agents may help overcome this.

Alfaxalone produces dose-dependent respiratory depression and apnea is common after rapid intravenous injection of clinically relevant doses (Muir et al. 2008, 2009). Slow injection over 60 seconds is recommended and the anesthetist should be prepared to intubate, provide supplemental oxygen, and support ventilation when administering alfaxalone. Cardiac output and arterial blood pressure remain generally stable after administration of clinically appropriate doses in healthy patients, although caution should be exercised when administering alfaxalone to patients with limited cardiac reserve capacity or hemodynamic instability.

In addition to dogs and cats, off-label alfaxalone use has been reported in many other domestic species and exotic pets, including horses, sheep, swine, rabbits, ferrets, rodents (excluding mice, rats, guinea pigs, chinchillas), birds (excluding budgerigars and finches), reptiles (excluding lizards, turtles, tortoises, and snakes), and amphibians (excluding frogs, toads, and axolotls). It has also been used in a diverse range of captive zoo species (excluding civet cats, macaques, sea lions, crocodiles, pythons, flamingos, parrots, blue crabs, carp, and tarantulas); as well as in free-ranging wildlife (excluding deer, bighorn sheep, seals, koalas, wallabies, and river turtles).

## Opioids

All drugs classified as opioids are chemically related to a group of compounds which have been purified from the juice of a particular species of poppy, *Papaverum somniferum*. The unrefined extract from the poppy is called opium and contains approximately 20 naturally occurring pharmacologically active compounds, including morphine and codeine. In addition, numerous semisynthetic and synthetic analogs of these natural compounds have been developed for clinical use. The word “opioid” is typically used to encompass all chemical derivatives of the compounds purified from opium and will be the term used to describe this class of analgesics throughout this section.

The opioids are a versatile group of drugs with extensive applications related to the management of pain in companion animal veterinary medicine. In the past, their use in wild and exotic species has been largely limited to the ultra-potent agents utilized in remote capture techniques. However, with the rapid evolution of zoo animal medicine and surgery, the opioids are being used increasingly as analgesics for the management of surgical pain in a wide variety of species. Although there are few pharmacokinetic or pharmacodynamic studies involving opioids in wild and exotic animals, a general discussion of opioid pharmacology is relevant and may facilitate extrapolation from companion animal species.

### Opioid Receptors

It is well known that exogenously administered opioids such as morphine or heroin exert their effects by interacting with specific opioid receptors and mimicking naturally occurring molecules known as endogenous opioid peptides. It is now accepted that there are three well-defined types of opioid receptors, most commonly known by their Greek letter designations as  $\mu$  (mu),  $\delta$  (delta), and  $\kappa$  (kappa) (Harrison et al. 1998; Kieffer 1999; Janecka et al. 2004; Inturrisi 2002). This classic system of nomenclature has been under reconsideration for a number of years and during this time several alternative naming systems have been proposed, leading to considerable confusion. In addition, a fourth type of opioid receptor, the nociceptin receptor (also known as the orphanin FQ receptor), has been characterized (Moran et al. 2000; Smith et al. 2001). According to the most recent recommendations of the International Union of Basic and Clinical Pharmacology Committee on Receptor Nomenclature and Drug Classification (NC-IUPHAR), variations based on the Greek letters remain acceptable. Thus, mu,  $\mu$ , or MOP (for “mu opioid peptide”); delta,  $\delta$  or DOP (for “delta opioid peptide”); kappa,  $\kappa$  or KOP (for “kappa opioid peptide”); and NOP (for “nociceptin opioid peptide”) are considered interchangeable abbreviations. Distinct cDNA sequences have been cloned for all four opioid receptor types, and each type appears to have a unique distribution in the brain, spinal cord, and periphery (Smith & Lee 2003).

The diversity of opioid receptors is further extended by the existence of several subtypes of  $\mu$ ,  $\delta$ , and  $\kappa$  receptors. Based on pharmacologic studies, there are thought to be at least three  $\mu$  receptor subtypes,  $\mu_1$ ,  $\mu_2$ , and  $\mu_3$ ; two  $\delta$  receptor subtypes,  $\delta_1$ , and  $\delta_2$ ; and perhaps as many as four  $\kappa$  receptor subtypes,  $\kappa_{1a}$ ,  $\kappa_{1b}$ ,  $\kappa_2$ , and  $\kappa_3$  (Smith & Lee 2003). The discovery of opioid receptor subtypes generated great enthusiasm among researchers and introduced the possibility of developing subtype-specific therapeutic agents

with favorable side-effect profiles. At this point, however, the functional significance of these receptor subtypes remains unclear and distinct cDNA sequences corresponding to these subtypes have not yet been identified (Smith & Lee 2003).

In general, it appears that the  $\mu$  receptor mediates most of the clinically relevant analgesic effects, as well as most of the adverse side effects associated with opioid administration (Kieffer 1999). Drugs acting at the  $\delta$  receptor tend to be poor analgesics but may modify  $\mu$  receptor-mediated antinociception under certain circumstances and mediate opioid receptor “cross-talk”. The  $\kappa$  receptor mediates analgesia in several specific locations in the CNS and the periphery, but distinguishing  $\mu$ - and  $\kappa$ -mediated analgesic effects has proven to be difficult (Kieffer 1999; Smith & Lee 2003). In contrast to the classic opioid receptors, the nociceptin receptor does not mediate typical opioid analgesia, but instead produces anti-opioid (pro-nociceptive) effects (Janecka et al. 2004; Moran et al. 2000; Smith et al. 2001). Due to the considerable structural homology between the three classically described opioid receptors, it is likely that there are significant interactions between these receptors in different tissues, and the loosely defined physiologic roles ascribed to each receptor type still require further clarification.

### Endogenous Opioid Receptor Ligands

The opioid receptors discussed above are part of an extensive opioid system that includes a large number of endogenous opioid peptide ligands. Endogenous opioid peptides are small molecules that are naturally produced in the CNS and in various glands throughout the body, such as the pituitary and the adrenal (Janecka et al. 2004). Three classical families of endogenous opioid peptides have been identified: the enkephalins, the dynorphins, and  $\beta$ -endorphin. Each of these is derived from a distinct precursor polypeptide, pro-enkephalin, pro-dynorphin, and pro-opiomelanocortin, respectively (Janecka et al. 2004). These classical endogenous opioid peptides are expressed throughout the CNS and their presence has more recently been confirmed in peripheral tissues as well (Janecka et al. 2004). There are considerable structural similarities between these three groups of peptides, and each family demonstrates variable affinities for  $\mu$ ,  $\delta$ , and  $\kappa$  receptors. None of them bind exclusively to a single opioid receptor and none of them have any significant affinity for the nociceptin receptor. The physiologic roles of these peptides are not completely understood at this time. They appear to function as neurotransmitters, neuromodulators and, in some cases, as neurohormones. They mediate some forms

of stress-induced analgesia and play a role in analgesia induced by electrical stimulation of discrete regions in the brain, such as the periaqueductal gray of the mesencephalon (Inturrisi 2002).

Nociceptin (also known as orphanin FQ) is the endogenous ligand for the more recently discovered nociception receptor. Nociceptin is derived from pro-nociceptin, and its amino acid sequence is closely related to that of the classical endogenous opioid peptides discussed above (Janecka et al. 2004; Moran et al. 2000). Despite this homology, nociception binding is specific for the nociception receptor and the peptide does not appear to interact with  $\mu$ ,  $\delta$ , or  $\kappa$  receptors. Furthermore, the physiologic effects of nociceptin are in direct contrast to the actions of the classical endogenous opioid peptides, with nociception producing a distinctly pro-nociceptive effect (Janecka et al. 2004; Moran et al. 2000; Smith 2001). The functional significance of nociception and its receptor remains to be elucidated, but additional insight into this novel opioid peptide may have substantial implications in future therapeutic drug development.

In addition to the enkephalins, dynorphins,  $\beta$ -endorphin, and nociception, there are now two other discovered endogenous opioid peptides called endomorphin-1 and endomorphin-2 (Zadina et al. 1990). These peptides are putative products of an, as yet, unidentified precursor and have been proposed to be the highly selective endogenous ligands for the  $\mu$  receptor (Janecka et al. 2004; Zadina et al. 1990). The endomorphins are small tetrapeptides that are structurally unrelated to the classical endogenous opioid peptides (Zadina et al. 1990). Their identification has heralded a new era in research of the  $\mu$  opioid system which may contribute to our understanding of the neurobiology of opioids, and provide new avenues for therapeutic interventions.

Another endogenous opioid receptor agonist, dermorphin, was discovered in the early 1980s in the skin of a South American frog species belonging to the genus *Phyllomedusa*. Dermorphins are full  $\mu$  receptor agonists with approximately 30–40 times the potency of morphine. Shortly after the discovery of amphibian dermorphin, another endogenous opioid, deltorphin, was isolated and shown to act as an agonist at  $\delta$  receptors. These natural opioid peptides appear to be unique to amphibians and, although their functional role is not entirely clear, it has been speculated that they may be involved in protection from predators and/or hibernation (Mosely & Mosely 2015). While the discovery of the dermorphins generated initial interest as potential clinical analgesic agents, little work has been done in recent years. Dermorphin has been used illegally in horse racing as a performance-enhancing drug (Robinson et al. 2015).

## Opioid Receptor Signaling and Mechanisms of Analgesia

Binding of an opioid agonist to a neuronal opioid receptor, regardless of whether the agonist is endogenous or exogenous, typically leads to several events that serve to inhibit the activation of the neuron. Opioid receptors are part of a large superfamily of membrane-bound receptors that are coupled to G-proteins (Smith & Lee 2003). As such, they are structurally and functionally related to receptors for many neurotransmitters and other neuropeptides which act to modulate the activity of nerve cells. Opioid receptor binding, via activation of various types of G-proteins, may lead to inhibition of adenylyl cyclase (cAMP) activity, activation of receptor-operated  $K^+$  currents, and suppression of voltage-gated  $Ca^{2+}$  currents (Inturrisi 2002).

At the pre-synaptic level, decreased  $Ca^{2+}$  influx will result in reduced release of transmitter substances, such as substance P, from primary afferent fibers in the spinal cord dorsal horn, thereby inhibiting synaptic transmission of nociceptive input (Inturrisi 2002). Post-synaptically, enhanced  $K^+$  efflux will result in neuronal hyperpolarization of spinal cord projection neurons and inhibition of ascending nociceptive pathways. A third potential mode of opioid action involves upregulation of supraspinal descending antinociceptive pathways in the periaqueductal gray. It is now known that this system is subject to tonic inhibition mediated by GABAergic neurons, and opioid receptor activation has been shown to suppress this inhibitory influence and augment descending antinociceptive transmission (Inturrisi 2002; Christie et al. 2000). The proposed cellular basis for this involves  $\mu$  receptors present on pre-synaptic GABAergic nerve terminals which activate voltage-dependent  $K^+$  currents and thereby inhibit GABA release into the synaptic cleft (Christie et al. 2000). It is important to note that while our collective understanding of opioid receptor-mediated signaling has increased dramatically in recent years, the relationship of such subcellular events to clinical analgesia at the level of the organism requires further clarification.

## Opioid Receptor Distribution and Therapeutic Implications

While cellular and molecular studies of opioid receptors and ligands are invaluable in understanding their function, it is critical to place them in their anatomical and physiologic context to fully appreciate the opioid system and its relevance to pain management. It has long been a principle tenet of opioid analgesia that these agents are centrally acting, and this understanding has shaped the way we use opioid analgesics clinically. It has been well

established that the analgesic effects of opioids arise from their ability to directly inhibit the ascending transmission of nociceptive information from the spinal cord dorsal horn, and to activate pain control circuits that descend from the midbrain via the rostral ventromedial medulla to the spinal cord. Within the CNS, evidence of  $\mu$ ,  $\delta$ , and  $\kappa$  opioid receptor mRNA and/or opioid peptide binding has been demonstrated in supraspinal sites, including the mesencephalic periaqueductal gray, the mesencephalic reticular formation, various nuclei of the rostral ventromedial medulla, forebrain regions including the nucleus accumbens, as well as spinally within the dorsal horn (Gutstein & Akil 2001; Yaksh et al. 1998). The interactions between groups of opioid receptors at various spinal and supraspinal locations, as well as interactions between different receptor types within a given location, are complex and incompletely understood at this time.

Systemic administration of opioid analgesics via intravenous, intramuscular, or subcutaneous injection will result in a relatively rapid onset of action via interaction with these CNS receptors. Oral, transdermal, rectal, or buccal mucosal administration of opioids will result in variable systemic absorption, depending on the characteristics of the particular agent, with analgesic effects being mediated largely by the same receptors within the CNS. In addition, neuraxial administration, into either the subarachnoid or epidural space, is a particularly efficacious route of administration. Small doses of opioids introduced via these routes readily penetrate the spinal cord and interact with spinal and/or supraspinal opioid receptors to produce profound and potentially long-lasting analgesia, the characteristics of which will depend on the particular drug utilized.

Despite the fact that opioids have long been considered the prototype of centrally acting analgesics, a body of evidence has emerged over the past decade which clearly indicates that opioids can produce potent and clinically measurable analgesia by activation of opioid receptors in the peripheral nervous system (Stein et al. 2001). Opioid receptors of all three major types have been identified on the processes of sensory neurons, and they respond to peripherally applied opioids and locally released endogenous opioid peptides when upregulated during inflammatory pain states (Stein 1993; Stein et al. 1993, 2001, 2003; Fields et al. 1980). Furthermore, sympathetic neurons and immune cells have also been shown to express opioid receptors, but their functional role remains unclear (Stein et al. 2003). While the binding characteristics of peripheral and central opioid receptors are similar, the molecular mass of peripheral and central  $\mu$  opioid receptors appears to be different, suggesting that selective ligands for these

peripheral receptors could be developed which would produce opioid analgesia without the potential to induce centrally mediated adverse side effects (Stein 1995, Stein & Yassouridis 1997; Stein et al. 1996, 2001, 2003).

## Adverse Effects of Clinically Used Opioids

While opioids are used clinically primarily for their pain-relieving properties, they also produce a host of other effects on a variety of body systems. This is not surprising in light of the wide distribution of endogenous opioid peptides and their receptors in supraspinal, spinal, and peripheral locations. Since information regarding opioid side effects in most wild and exotic animals is lacking, reference is made to common domestic species where appropriate.

### Central Nervous System

#### Level of Arousal

There are considerable species differences in the CNS response to opioid analgesics that cannot be attributed to pharmacokinetic variations alone. CNS depression (i.e., sedation) is typically seen in the dog, monkey, and human, while CNS stimulation (i.e., excitement and/or spontaneous locomotor activity) may be elicited in the cat, horse, goat, sheep, pig, and cow following systemic administration of various opioids, most notably morphine (Branson et al. 2001). The reasons for these different responses are not entirely clear at this time, but are presumably related to differing concentrations and distributions of  $\mu$ ,  $\delta$  and  $\kappa$  receptors in various regions of the brain in these species (Hellyer et al. 2003). Details regarding the CNS responses of specific wild and exotic species to opioids are not yet known. Regardless of the species, however, there are numerous factors which may play a role, including the temperament or condition of the animal, the presence or absence of pain in the animal, the dose, route and timing of drug administration, and the specific opioid administered.

#### Thermoregulatory Center

The hypothalamic thermoregulatory system is also affected by opioid administration. Hypothermia tends to be the most common response, particularly when opioids are used during the perioperative period in the presence of other CNS depressant drugs (Gutstein & Akil 2001; Branson et al. 2001). Under some clinical circumstances, however, hyperthermia is observed in cats, horses, swine, and ruminants following opioid administration (Niedfeldt & Robertson 2006; Posner et al. 2007, 2010). Part of this increase in body temperature may be attributed to an increase in muscle activity associated with CNS excitation in these species, but a specific central hypothalamic mechanism has also been implicated, although it remains poorly understood (Branson et al. 2001).

### **Emetic Center**

Nausea and vomiting associated with opioid administration are a result of direct stimulation of the chemoreceptor trigger zone for emesis located in the area postrema of the medulla (Gutstein & Akil 2001; Stoelting 1999). As with the other centrally mediated side effects, species plays a role in determining an individual's tendency to vomit after an opioid is administered. Horses, rabbits, ruminants, and swine do not vomit with opioid administration. Cats may vomit, but usually at doses that are greater than those which stimulate vomiting in dogs. Dogs will commonly vomit following opioid administration, especially with morphine.

### **Cough Center**

Opioids have variable efficacy in depressing the cough reflex, at least in part, by a direct effect on a cough center located in the medulla (Gutstein & Akil 2001). Certain opioids are more effective antitussives than others, including drugs like codeine, hydrocodone, and butorphanol.

### **Pupillary Diameter**

As a general rule, opioids tend to produce mydriasis in those species that exhibit CNS excitation, and miosis in those that become sedated following opioid administration (Branson et al. 2001; Stephan et al. 2003; Lee & Wang 1975; Wallenstein & Wang 1979). Miosis is due to an excitatory action of opioids on neuronal firing in the oculomotor nucleus (Stoelting 1999; Lee & Wang 1975; Wallenstein & Wang 1979). In the cat, and presumably in other species that exhibit mydriasis, this increase in activity in the oculomotor nuclear complex still occurs but the miotic effect is masked by increased release of catecholamines, which results in mydriasis (Wallenstein & Wang 1979).

### **Respiratory System**

Opioids produce dose-dependent depression of ventilation, primarily mediated by  $\mu_2$  receptors leading to a direct depressant effect on brainstem respiratory centers (Gutstein & Akil 2001; Stoelting 1999). This effect is characterized by decreased responsiveness of these centers to carbon dioxide and is reflected in an increased resting  $\text{PaCO}_2$  and displacement of the carbon dioxide response curve to the right. This effect is compounded by the coadministration of sedative and/or anesthetic agents, meaning that significant respiratory depression and hypercapnia are much more likely to occur in anesthetized patients who receive opioids versus those who are conscious. It should be noted that, in general, humans tend to be more sensitive to the respiratory depressant effects of opioids when compared to most veterinary species. However, respiratory support and/or specific opioid antagonists should be immediately available anytime very high doses, or very potent opioids are used.

### **Cardiovascular System**

Most opioids have minimal effects on cardiac output, cardiac rhythm, and arterial blood pressure when clinically relevant analgesic doses are administered. Bradycardia may occur as a result of opioid-induced medullary vagal stimulation and will respond readily to anticholinergic treatment if warranted. Particular opioids (morphine and meperidine) can cause release of histamine, especially after rapid intravenous administration, which may lead to vasodilation and hypotension (Branson et al. 2001; Smith et al. 2001). Due to their relatively benign effects on cardiovascular function, opioids commonly form the basis of anesthetic protocols for human patients or animals with preexisting cardiovascular disease.

### **Gastrointestinal System**

The gastrointestinal effects of the opioids are mediated by  $\mu$  and  $\delta$  receptors located in the myenteric plexus of the gastrointestinal tract (Gutstein & Akil 2001; Branson et al. 2001). Opioid administration may stimulate defecation in certain species. Following this initial response, spasm of gastrointestinal smooth muscle predisposes to ileus and constipation. Horses and ruminants in particular may be predisposed to gastrointestinal complications associated with opioid administration such as colic and ruminal tympany, respectively. Chronic opioid use may predispose to gastrointestinal stasis in other species.

In human patients, opioids (most notably fentanyl and morphine) have been shown to increase bile duct pressure through constriction of the sphincter of Oddi (Radnay et al. 1984). The incidence of this side effect in humans is, however, quite low (Jones et al. 1981). The incidence of increased bile duct pressure secondary to opioid administration in various animal species, and its potential clinical significance, is unknown at this time.

### **Genitourinary System**

Opioids, particularly when administered neuraxially, may cause urinary retention through dose-dependent suppression of detrusor contractility and decreased sensation of urge (Kuipers et al. 2004; el Bindary & Abu el-Nasr 2001). Urine volume may also be affected by opioids and the mechanism of this effect appears to be multifactorial.  $\mu$  agonists tend to produce oliguria in the clinical setting and this is in part a result of increased antidiuretic hormone release leading to altered renal tubular function (Stoelting 1999; Mercadante & Arcuri 2004). Elevations in circulating plasma atrial natriuretic peptide may also play a role in morphine-induced antidiuresis (Mercadante & Arcuri 2004). Conversely,  $\kappa$  agonists tend to produce a diuretic effect, possibly through inhibition of antidiuretic hormone secretion (Stoelting 1999; Mercadante & Arcuri 2004).

Other peripheral mechanisms involving stimulation of renal alpha-2 adrenergic receptors may also contribute to this  $\kappa$  agonist effect (Mercadante & Arcuri 2004).

### Opioid-Induced Hyperalgesia


There is a growing body of evidence from both human and rodent studies supporting the existence of a condition called opioid-induced hyperalgesia (Lee et al. 2011; Gomez et al. 2009; Yu et al. 2016; Colvin et al. 2019). This phenomenon is characterized by a paradoxical increase in pain associated with opioid administration and can contribute to poorly controlled pain and dose escalation. In humans, it has been observed with acute and chronic exposure, high and low doses, and with different types of opioids and routes of administration. The mechanisms underlying opioid-induced hyperalgesia are not well understood but both  $\mu$  opioid receptors on transient receptor potential vanilloid-1 (TRPV1)-expressing dorsal root ganglion neurons and NMDA receptors in the spinal cord dorsal horn appear to be involved (Deng et al. 2019). At this time, animal studies involving opioid-induced hyperalgesia are largely confined to experimental pain models in rodents, however, awareness in veterinary medicine is increasing and publications involving dogs and cats are beginning to emerge (Ruíz-López et al. 2020; Borland & Garcia 2019; White et al. 2017).

### Opioid Agonists

Almost all clinically useful opioids exert their analgesic and immobilizing effects by acting as agonists at  $\mu$  receptors. While there are a few opioids which act as  $\kappa$  agonists, these drugs also tend to have antagonist or partial agonist effects at  $\mu$  and/or  $\delta$  receptors and are thus not classified as *pure* agonists. Pure or full opioid agonists are capable of eliciting maximal activation of the receptor when they bind it and the subsequent downstream processes result in a maximal analgesic effect (Fig. 1.1). Clinically, the full  $\mu$  agonists are superior analgesics and they are the drugs of choice for pain of moderate to severe intensity in many veterinary species. The following section contains brief descriptions of full  $\mu$  agonists currently used in veterinary medicine. Specific details regarding  $\mu$  agonist clinical pharmacology in various wild and exotic species are lacking.

#### Morphine (Morphine Sulfate)

Morphine is the prototypical opioid analgesic and acts as a full agonist not only at  $\mu$  receptors, but also at  $\delta$  and  $\kappa$  receptors (Gutstein & Akil 2001). Despite the development of numerous synthetic opioids, many of which are more potent than morphine and may have other characteristics that make them desirable alternatives to morphine in certain



**Figure 1.1** Effects of opioid agonists.

circumstances, it is worth noting that no other drug has been shown to be more efficacious than morphine at relieving pain in humans. Compared to the synthetic opioid agonists, morphine is relatively hydrophilic in nature and crosses the blood-brain barrier more slowly than fentanyl or oxycodone, thereby delaying the peak effect somewhat even after intravenous administration (Gutstein & Akil 2001; Stoelting 1999). Clinically, this lag is not likely to be significant under most circumstances, with the onset of analgesia occurring reasonably promptly after a single dose of morphine and typically lasting 3–4 hours (Barnhart et al. 2000; Taylor et al. 2001). Morphine's poor lipid solubility means that it can produce long-lasting analgesia when administered into the epidural or subarachnoid space, with effects persisting for 12–24 hours. The first-pass effect is significant after oral administration and the bioavailability of oral morphine preparations is at best only in the range of 0–25%.

In most species, the primary metabolic pathway for morphine involves conjugation with glucuronic acid leading to the formation of two major metabolites, morphine-6-glucuronide and morphine-3-glucuronide (Gutstein & Akil 2001; Faura et al. 1998). Despite the low levels of glucuronyl transferase in the cat, the pharmacokinetics of morphine in this species seem to be broadly comparable to the dog and human, though clearance rates may be marginally slower (Barnhart et al. 2000; Taylor et al. 2001; Faura et al. 1998). This suggests that morphine must undergo a different type of conjugation reaction in this species. Morphine-6-glucuronide

has pharmacological activities that are indistinguishable from those of morphine in animal models and in human beings, while morphine-3-glucuronide appears to have little affinity for opioid receptors, but may contribute to the excitatory effects of morphine in some situations (Gutstein & Akil 2001; Smith 2000). With chronic morphine administration, it is likely that the active metabolite, morphine-6-glucuronide, contributes significantly to clinical analgesia.

Very little morphine is excreted unchanged in the urine. The major metabolites, morphine-3-glucuronide and, to a lesser extent, morphine-6-glucuronide, are eliminated almost entirely via glomerular filtration. In human patients, renal failure may lead to accumulation of morphine-6-glucuronide and persistent clinical effects, while liver dysfunction seems to have minimal impact on morphine clearance (Gutstein & Akil 2001; Stoelting 1999).

The side effects associated with morphine administration are typical of most opioid agonists and have been discussed previously in this chapter. In particular, the increased incidence of vomiting after morphine administration as well as its potential to cause histamine release after intravenous administration distinguish morphine from other full opioid agonists.

### **Hydromorphone**

Hydromorphone is a synthetic opioid that acts as a full agonist at  $\mu$  receptors and is used in both human and veterinary medicine. Clinically, hydromorphone and oxymorphone have similar efficacy, potency, duration of analgesic action, and side-effect profiles, but hydromorphone remains significantly less expensive. Like oxymorphone, hydromorphone is not associated with histamine release so bolus intravenous administration is considered safe (Smith et al. 2001).

### **Meperidine**

Meperidine is a synthetic opioid that exerts its analgesic effects through agonism at  $\mu$  receptors. Interestingly, it also appears able to bind other types of receptors, which may contribute to some of its clinical effects other than analgesia. Meperidine is capable of blocking sodium channels and inhibiting activity in dorsal horn neurons in a manner analogous to local anesthetics (Wagner et al. 1999; Wolff et al. 2004). It has also recently been shown that meperidine exerts agonist activity at  $\alpha$ -2 receptors, specifically the  $\alpha$ -2B subtype, suggesting that it may possess some  $\alpha$ -2 agonist-like properties (Takada et al. 1999, 2002).

Meperidine has a considerably shorter duration of analgesic action compared to morphine, oxymorphone or hydromorphone, typically not extending beyond 1 hour (Branson et al. 2001). Metabolic pathways vary among different species but, in general, most of the drug is

demethylated to normeperidine in the liver and then undergoes further hydrolysis and ultimately renal excretion (Branson et al. 2001; Taylor et al. 2001; Yeh et al. 1981). Normeperidine is an active metabolite and possesses approximately one half of the analgesic efficacy of meperidine (Gutstein & Akil 2001; Branson et al. 2001). Normeperidine has produced toxic neurologic side effects in human patients receiving meperidine for prolonged periods of time, especially in the presence of impaired renal function (Stoelting 1999; Stone et al. 1993).

Unlike most of the other opioids in clinical use, meperidine has been shown to produce significant negative inotropic effects when administered alone to conscious dogs (Priano & Vatner 1981). In addition, due to its modest atropine-like effects, meperidine tends to increase heart rate rather than predispose to bradycardia as is often seen with other opioids (Branson et al. 2001; Stoelting 1999). The clinical significance of these cardiovascular effects in the peri-anesthetic period has never been clearly ascertained. Like morphine, meperidine also causes histamine release when administered intravenously (Branson et al. 2001).

A rare but life-threatening drug interaction has been reported in human patients receiving meperidine that may have relevance in veterinary medicine. The combination of meperidine (and perhaps other opioids) with a monoamine oxidase inhibitor may lead to "serotonin syndrome" which is characterized by a constellation of symptoms including confusion, fever, shivering, diaphoresis, ataxia, hyperreflexia, myoclonus, and diarrhea (Sporer 1995; Bowdle 1998; Heinonen & Myllyla 1998; Tissot 2003). A monoamine oxidase inhibitor, selegiline (Deprenyl®), has been used in dogs to treat pituitary-dependent hyperadrenocorticism or to modify behaviour in patients with canine cognitive dysfunction. Though there have not, to date, been any reports of adverse meperidine-selegiline interactions in dogs, the veterinarian must be aware of the potential for complications if analgesia is required in patients receiving monoamine oxidase inhibitors. A recent study has evaluated the effects of other opioids (oxymorphone and butorphanol) in selegiline-treated dogs and did not identify any specific adverse drug interactions in these animals (Dodam et al. 2004).

### **Fentanyl**

Fentanyl is a highly lipid-soluble, short-acting synthetic  $\mu$  opioid agonist. A single dose of fentanyl administered intravenously has a more rapid onset and a shorter duration of action than morphine. Peak analgesic effects occur in about 5 minutes and last approximately 30 minutes (Gutstein & Akil 2001; Stoelting 1999). Rapid redistribution of the drug to inactive tissue sites, such as fat and skeletal

muscle, leads to a decrease in plasma concentration and is responsible for the prompt termination of clinical effects. In most veterinary species the elimination half-time after a single bolus or a brief infusion is in the range of 2–3 hours (Carroll et al. 1999; Lee et al. 2000; Maxwell et al. 2003). Administration of very large doses or prolonged infusions may result in saturation of inactive tissues with termination of clinical effects becoming dependent on hepatic metabolism and renal excretion (Gutstein & Akil 2001; Stoelting 1999). Thus, the context-sensitive half-time of fentanyl increases significantly with the duration of the infusion, and clinical effects may persist for an extended period following termination of a long-term intravenous infusion.

Side effects associated with fentanyl administration are similar to those of the other full  $\mu$  agonist opioids. In general, cardiovascular stability is excellent with fentanyl and intravenous administration is not associated with histamine release (Gutstein & Akil 2001; Stoelting 1999). Bradycardia may be significant with bolus doses but readily responds to anticholinergics if treatment is warranted (Gutstein & Akil 2001; Branson et al. 2001). In human patients muscle rigidity, especially of the chest wall, has been noted after administration of fentanyl or one of its congeners. The potential significance of this adverse effect in animal patients is not clear at this time, and the risk is considered minimal if large rapid bolus administrations are avoided.

Clinically, fentanyl is used most frequently in dogs and cats but it is also a potentially useful analgesic in other species, including the horse, cow, sheep, goat, and pig. Historically, fentanyl was available in combination with the butyrophenone tranquilizer droperidol in a product called Innovar-Vet, which was typically administered in the preanesthetic period to provide sedation and analgesia. This product is no longer available and systemic administration of fentanyl today is usually via the intravenous route.

The development of novel less-invasive routes of opioid administration for use in human patients led to the marketing of transdermal fentanyl patches (Duragesic®). The patches are designed to release a constant amount of fentanyl per hour which is then absorbed across the skin and taken up systemically. Fentanyl patches are designed for human skin and body temperature; however, their use has been evaluated in a number of domestic veterinary species (Carroll et al. 1999; Maxwell et al. 2003; Gilberto et al. 2003; Franks et al. 2000; Wilkinson et al. 2001; Egger et al. 1998, 2003; Gellaseh et al. 2002; Robinson et al. 1999). Substantial variations in plasma drug concentrations have been documented and significant lag times after patch placement are common prior to onset of analgesia (Carroll

et al. 1999; Lee et al. 2000; Egger et al. 1998, 2003). Furthermore, changes in body temperature have been shown to significantly affect fentanyl absorption in anesthetized cats, and it is likely that other factors associated with skin preparation and patch placement have the potential to substantially alter plasma fentanyl levels and analgesic efficacy (Pettifer & Hosgood 2003). Fentanyl patch safety and efficacy in most species is unknown at this time.

### **Alfentanil, Sufentanil, and Remifentanil**

Alfentanil, sufentanil, and remifentanil are all structural analogues of fentanyl that were developed for use in humans in an effort to create analgesics with a more rapid onset of action and predictable termination of opioid effects. All three are similar with regard to onset and all have context-sensitive half-times that are shorter than that of fentanyl after prolonged infusions (Stoelting 1999). Remifentanil is unique among opioids because it is metabolized by nonspecific plasma esterases to inactive metabolites (Chism & Rickert 1996; Hoke et al. 1997). Thus, hepatic or renal dysfunction will have little impact on drug clearance and this, in combination with the robust nature of the esterase metabolic system, contributes to the predictability associated with infusion of remifentanil (Gutstein & Akil 2001; Stoelting 1999).

All three of these drugs are used during general anesthesia for procedures requiring intense analgesia and/or blunting of the sympathetic nervous system response to noxious stimulation. These agents, specifically remifentanil, are the opioids most frequently associated with opioid-induced hyperalgesia in humans (Santonocito et al. 2018). As yet, they have limited applications for postoperative or chronic pain management. Like fentanyl, they can be administered at relatively low infusion rates as adjuncts to general anesthetic protocols based on volatile inhalant or other injectable agents, or they can be administered at higher rates as the primary agent for total intravenous anesthesia. The minimum alveolar anesthetic concentration-sparing properties of these agents have been demonstrated in both the dog and cat (Hoke et al. 1997; Michelsen et al. 1996; Mendes & Selmi 2004; Ilkiw et al. 1997; Pascoe et al. 1997). In the horse, systemic infusions of alfentanil did not have significant effects on minimum alveolar concentrations of inhalant anesthetics and when administered to conscious horses were associated with increases in locomotor activity (Pascoe et al. 1991, 1993; Pascoe & Taylor 2003). There is little evidence to suggest that any of the fentanyl analogues offer advantages over morphine when administered into the epidural space for analgesia (Natalini & Robinson 2000).

### Methadone

Methadone is a synthetic  $\mu$  opioid agonist with pharmacologic properties qualitatively similar to those of morphine, but possessing additional affinity for NMDA receptors (Gorman et al. 1997; Ripamonti & Dickerson 2001). Methadone's unique clinical characteristics include excellent absorption following oral administration, no known active metabolites, high potency, and an extended duration of action (Gutstein & Akil 2001; Branson et al. 2001; Ripamonti & Dickerson 2001). In human patients, the drug has been used primarily in the treatment of opioid abstinence syndromes but it is being used increasingly for the management of chronic pain. In contrast to humans, methadone has low bioavailability in most veterinary species and consequently has limited efficacy when administered orally. A racemic injectable formulation of methadone hydrochloride (Comfortan®) is approved for use in dogs and cats for analgesia and premedication prior to general anesthesia in Europe and the United Kingdom. Numerous publications involving intramuscular, intravenous, and subcutaneous methadone in these and other domestic species have emerged in recent years as the drug gains popularity in clinical veterinary practice (Bini et al. 2018; Bitti et al. 2017; Hunt et al. 2013; Slingsby et al. 2015; Shah et al. 2018; Trimble et al. 2018; Warne et al. 2013; Gozalo-Marcilla et al. 2019).

### Codeine

Codeine is the result of substitution of a methyl group onto morphine which acts to limit first-pass hepatic metabolism and accounts for codeine's high oral bioavailability (Gutstein & Akil 2001; Stoelting 1999). Codeine is well known for its excellent antitussive properties and is often combined in an oral formulation with a nonopioid analgesic, such as acetaminophen (Tylenol 3®), for the management of mild to moderate pain in human patients.

### Oxycodone and Hydrocodone

Oxycodone and hydrocodone are opioids that are typically administered orally for the treatment of pain in human patients. Though oxycodone is available as a single drug continuous-release formulation (Oxycontin®), these drugs are most often prepared in combination with nonopioid analgesics such as aspirin and acetaminophen (Percocet®, Percodan®, Lorcet®, Vicodan®, etc.). In light of the epidemic of opioid addiction in humans and the potential for drug diversion, veterinarians are advised to prescribe alternative analgesics and avoid off-label use of opioids such as oxycodone and hydrocodone due to their extremely high abuse potential.

### Etorphine and Carfentanil (M-99® and Wildnil®, Respectively)


These two opioids are discussed together because they are both used exclusively for the restraint and capture of wild animals, rather than as analgesic agents. They are extremely potent opioids and the immediate availability of a suitable antagonist is mandatory before these drugs are to be used, not only to reverse drug effects in animal patients, but also as a safety precaution in the event of accidental human injection. Though etorphine and carfentanil are most often injected intramuscularly (usually using a remote drug delivery technique) recent studies suggest that carfentanil is useful when administered orally in a variety of species, including the brown bear, the Brazilian tapir, and the chimpanzee (Mortenson & Bechert 2001; Pollock & Ramsay 2003; Kearns et al. 2000; Mama et al. 2000). A number of different drugs have been used in combination with etorphine or carfentanil to enhance muscle relaxation, including acepromazine, xylazine, and medetomidine (Caulkett et al. 2000; Miller et al. 2003; Ramdohr et al. 2001; Roffe et al. 2001).

### Thiafentanil


Thiafentanil is an opioid agonist that has been utilized to facilitate the capture of many species of mammals, mainly ungulates, as well as large birds. It is pharmacologically classified as a synthetic opioid that has a relatively short duration of action. The shorter duration of action in combination with its reversibility with the opioid antagonist naltrexone make it an attractive agent when long periods of narcotization are not desirable.

### Opioid Agonist-Antagonists and Partial Agonists

This group includes drugs that have varying opioid receptor binding profiles, but which have one thing in common: they all occupy  $\mu$  opioid receptors, but do not initiate a maximal clinical response. Drugs such as butorphanol and nalbuphine are classified as agonist-antagonists. They are competitive  $\mu$  receptor antagonists, but exert their analgesic actions by acting as agonists at  $\kappa$  receptors (Fig. 1.2). Buprenorphine, on the other hand, is classified as a partial agonist and binds  $\mu$  receptors but produces only a limited clinical effect (Fig. 1.3). These mixed agonist-antagonist drugs were developed for the human market in an attempt to create analgesics with less respiratory depression and addictive potential. Due to their opioid receptor binding affinities, the side effects associated with these drugs demonstrate a so-called "ceiling effect" whereby increasing doses do not produce additional adverse responses. Unfortunately, the benefits of this ceiling effect on ventilatory depression come at the expense of limited analgesic efficacy and only a modest ability to decrease anesthetic requirements.



**Figure 1.2** Activation of  $\kappa$  receptor, but occupation without action at the  $\mu$  receptor.



**Figure 1.3** Partial activation of  $\mu$  receptor.

The coadministration of opioids with differing receptor binding profiles is currently an active area of research that deserves further attention. The interactions in this setting are complex and it appears that opioid coadministration has the potential to produce additive, synergistic, or antagonistic analgesic effects depending on the particular species, dosage, drugs, and pain model being evaluated. The following section contains brief descriptions of opioid

agonist-antagonists and partial agonists that are currently used in veterinary medicine.

### Butorphanol

Butorphanol is a synthetic agonist-antagonist opioid and has been used extensively in a wide variety of veterinary species. The drug was originally labelled as an antitussive agent in dogs and, even now, is approved as an analgesic in the cat and horse only (Branson et al. 2001). Butorphanol exerts its relevant clinical effects through its interactions at  $\kappa$  receptors and acts as an antagonist at  $\mu$  receptors. The duration of butorphanol's analgesic effects remains somewhat debatable and likely varies with species, type and intensity of pain, dosage, and route of administration (Sawyer et al. 1991; Robertson et al. 2003a; Sellon et al. 2001). In general, its effects are shorter-lived than those of morphine and are probably in the range of one to three hours. Butorphanol does not induce histamine release when administered intravenously and has minimal effects on cardiopulmonary function. There is conflicting evidence regarding the effects of butorphanol on inhalant anesthetic requirements in the dog, cat, and horse. Earlier studies failed to demonstrate a significant sparing effect on minimum alveolar concentration when butorphanol was coadministered with halothane in dogs and ponies (Quandt et al. 1994; Doherty et al. 1997; Matthews & Lindsay 1990). More recently, isoflurane MAC reductions have been documented after administration of clinically relevant doses of butorphanol in both dogs and cats (Ilkiw et al. 2002; Ko et al. 2000). The reasons for these discrepancies are probably related to differences in study techniques and, in the dog and cat specifically, it seems that butorphanol is capable of inducing at least modest reductions in inhalant anesthetic requirements.

Traditionally, it was thought that the simultaneous or sequential administration of butorphanol with a pure  $\mu$  opioid agonist such as morphine or hydromorphone would be counterproductive from an analgesic standpoint because butorphanol's ability to antagonize  $\mu$  receptors could inhibit or even reverse the effects of the agonist drug. Certainly, it has been clearly demonstrated that excessive sedation associated with a pure  $\mu$  agonist can be partially reversed by the administration of low doses of butorphanol, and it was presumed that butorphanol would similarly reverse the  $\mu$ -mediated analgesic effects as well. It would now appear that the potential interactions between butorphanol and full  $\mu$ -opioid agonists are more complex than originally believed. One study demonstrated that coadministration of butorphanol and oxymorphone to cats subjected to a visceral noxious stimulus resulted in enhanced analgesic effects (Briggs et al. 1998). A more recent feline study, however, evaluated the combination of butorphanol

and hydromorphone in a thermal threshold pain model and failed to demonstrate enhanced analgesia and suggested that butorphanol did, in fact, inhibit hydromorphone's analgesic effects (Lascelles & Robertson 2004). These contradictory findings illustrate that we still have much to learn about coadministration of opioid agents with differing receptor binding profiles and the clinical effects produced by such coadministration likely depend on many factors, including species, type of pain, dose, and the specific drugs involved.

### Nalbuphine and Pentazocine

Nalbuphine and pentazocine are classified as agonist-antagonist opioids and are clinically similar to butorphanol. They induce mild analgesia accompanied by minimal sedation, respiratory depression, or adverse cardiovascular effects. Like butorphanol, nalbuphine is occasionally used to partially reverse the effects of a full  $\mu$  agonist opioid while maintaining some residual analgesia.

### Buprenorphine

Buprenorphine is a semisynthetic, highly lipophilic opioid derived from thebaine. Unlike other opioids in this category, buprenorphine is considered to be a partial agonist at  $\mu$  opioid receptors. The drug binds avidly to and dissociates slowly from  $\mu$  receptors, but is not capable of eliciting a maximal clinical response. Due to its receptor binding characteristics, buprenorphine has a delayed onset of action and takes at least 60 minutes to attain peak effect after intramuscular administration. It also has a relatively long duration of action with clinical analgesic effects persisting for 6–12 hours in most species. Also, its high affinity for the  $\mu$  receptor means that it may be difficult to antagonize its effects with a drug such as naloxone. Buprenorphine has most often been administered intravenously or intramuscularly, however, due to the long lag time before clinical effects are achieved after intramuscular administration, the intravenous route is preferred. Comparable plasma drug levels and analgesic efficacy with oral transmucosal administration versus intravenous administration has been demonstrated in cats (Robertson et al. 2003). Compounded versions of buprenorphine are widely available and include higher concentrations for convenient dosing to larger animals and sustained release formulations for increased duration of action. Reports describing use of the compounded sustained-release buprenorphine formulation by ZooPharm are available in a wide range of zoo and wildlife species. If sustained release preparations are used, there should be a plan for supportive care should significant adverse effects occur since complete reversal of buprenorphine with conventional antagonists is often unsuccessful. In addition, a commercially available highly concentrated formulation (1.8 mg/mL) called Simbadol® has been


approved for perioperative analgesia in domestic cats with a label dose of 0.24 mg/kg administered subcutaneously every 24 hours for up to 3 days. Extralable intravenous and intramuscular administration has been used at lower doses and dosing intervals comparable to traditional buprenorphine doses. The duration of action of Simbadol is not due to any specific formulation characteristics. Rather it is a function of administration into adipose tissue which results in a gradual release into the circulation. A recent report describes the pharmacokinetics of highly concentrated buprenorphine/Simbadol® in Rhesus macaques (Mackiewicz et al. 2019).

## Opioid Antagonists

These drugs have high affinities for the opioid receptors and are able to displace opioid agonists from  $\mu$  and  $\kappa$  receptors. After this displacement, the pure antagonists bind to and occupy opioid receptors, but do not activate them (Fig. 1.4). Under ordinary circumstances, in patients that have not received exogenous agonist opioids, the opioid antagonists have few clinical effects when administered at clinically relevant dosages (Gutstein & Akil 2001). It is important to recognize that these drugs will rapidly reverse all opioid-induced clinical effects, including analgesia.

### Naloxone

This pure opioid antagonist is capable of reversing all opioid agonist effects, producing increased alertness, responsiveness, coordination, and, potentially, increased perception of pain. Naloxone's duration of action is shorter than many



**Figure 1.4** No activation of receptors.

of the opioid agonists with recommended intravenous doses lasting between 30 and 60 minutes. Consequently, animals need to be closely monitored for renarcotization after a dose of naloxone. Occasionally, excitement or anxiety may be seen after naloxone reversal of an opioid agonist. Premature ventricular contractions have also been documented after reversal, but are not a common occurrence and seem to be more likely if there are high levels of circulating catecholamines. This drug is sometimes administered sublingually to neonatal patients exhibiting respiratory depression that have been delivered by Caesarean section after maternal administration of an opioid agonist.

#### Nalmefene and Naltrexone

Both of these drugs are pure opioid antagonists with clinical effects that last approximately twice as long as naloxone (Veng-Pedersen et al. 1995). Though little is published about the use of these drugs in veterinary patients, they may be advantageous in preventing renarcotization when used to antagonize the effects of a long-acting opioid.


## Nonsteroidal Anti-inflammatories

The nonsteroidal anti-inflammatory analgesics (NSAIDs) relieve mild to moderately severe pain and have been used extensively in a wide variety of domestic animals for many years. While pharmacokinetic and pharmacodynamic studies involving NSAIDs in wild and exotic animals are lacking, their use in such species continues to increase. This class of drugs dates back thousands of years, with the salicylates being among the oldest and still most commonly used analgesics (Vane & Botting 2003). Salicylate is a naturally occurring substance found in willow bark and was used for centuries to manage pain associated with rheumatism prior to production of the synthetic compound. In 1878, Felix Hoffman working at the Bayer company in Germany made the acetylated form of salicylic acid which has come to be known as aspirin (Vane & Botting 2003). While aspirin (acetylsalicylic acid [ASA]) has been found to be effective in the management of acute and chronic mild discomfort, the newer injectable NSAIDs appear to have comparable efficacy to the pure  $\mu$  agonist opioids in controlling moderate to severe soft tissue and orthopedic pain. The NSAIDs appear to confer synergism when used in combination with opioids and may demonstrate an opioid-sparing effect should lower dosages of opioid be required. Their extended duration of action, in addition to their analgesic efficacy, make the NSAIDs ideal for treating acute and chronic pain in veterinary species. Careful patient and drug selection is critical, however, due to their potential for harmful side effects.

## Cyclooxygenases and Prostaglandin Synthesis

In 1971, Vane discovered the mechanism by which aspirin exerts its anti-inflammatory, analgesic, and antipyretic actions. He proved that aspirin and other NSAIDs inhibited the activity of a cyclooxygenase (COX) enzyme which produced prostaglandins (PGs) involved in the pathogenesis of inflammation, swelling, pain, and fever (Vane 1971). Twenty years later, the discovery of a second COX enzyme was made and more recently COX-3 (Botting 2000, 2003; Chandrasekharan et al. 2002). Cyclooxygenase (previously termed prostaglandin synthase) oxidizes arachidonic acid (previously termed eicosatetraenoic acid) to various eicosanoids (including PGs and other related compounds) (Fig. 1.5) (Livingston 2000). Oxidation of arachidonic acid by 5-lipoxygenase (5-LOX), the most biologically important of the mammalian oxygenases, results in the series of eicosanoids termed leukotrienes (Fig. 1.5). The release of arachidonic acid from membrane phospholipid is catalyzed by the enzyme phospholipase  $A_2$  and is the rate-limiting step in PG and leukotriene synthesis. Prostaglandin  $G_2$  is the initial prostanoid formed, followed by prostaglandin  $H_2$  which serves as a substrate for prostaglandin E-synthetase, prostaglandin D-isomerase, prostaglandin F-reductase, prostacyclin synthetase, and thromboxane synthetase for conversion to a variety of other prostenoids ubiquitous throughout cells and tissues in the body (Livingston 2000). These include the PGs  $PGE_2$ ,  $PGD_2$ ,  $PGF_2$ , and  $PGI_2$  (prostacyclin), and the thromboxanes  $TXA_2$  and  $TXB_2$ , all with diverse functions (Vane & Botting 1995). The PGs are not stored but are synthesized at a constant rate. They have short half-lives of 4–6 minutes at 37°C, and act locally at the site of production.

The PGs produced by both COX-1 and COX-2 are ubiquitous throughout the body and serve to facilitate many normal physiologic functions during both health and illness. Consequently, the clinical use of NSAIDs has the potential to disrupt these functions with the possibility of significant organ dysfunction. Thus, in addition to their role as analgesics, the effects of NSAIDs on the constitutive functions of the PGs must always be considered. There are several key points to note: (i) COX-1 generates PGs that are responsible for “mucosal defence” (i.e., secretion of bicarbonate and mucus, mucosal blood vessel attenuation of constriction, and mucosal epithelial regeneration), as well as thromboxane  $A_2$ , which is necessary for platelet function; (ii) COX-2 produces PGs which function in the prevention and promotion of healing of mucosal erosions, exert anti-inflammatory effects by inhibiting leukocyte adherence, and play a role in renal protection and maturation; and (iii) COX-3 produces PGs which exert a protective function by initiating fever (Vane & Botting 1995; Botting 2003).



**Figure 1.5** Eicosanoid synthesis.

Thus, depending on the NSAID selected, primary plug formation of platelets, modulation of vascular tone in the kidney and gastric mucosa, cytoprotective functions within the gastric mucosa, smooth muscle contraction, and regulation of body temperature will all be affected (Vane & Botting 1995). However, in this regard not all NSAIDs are created equal. As noted above, the COX-1, COX-2 and COX-3 enzymes make variable contributions to these functions, and individual NSAIDs inhibit each of these enzymes differently. Some NSAIDs inhibit both COX-1 and COX-2 (i.e., aspirin, phenylbutazone, ketoprofen/Anafen®, ketorolac/Toradol®, flunixin meglumine/Banamine®), while other NSAIDs preferentially inhibit COX-2 with only weak inhibition of COX-1 (i.e., meloxicam/Metacam®, carprofen/Rimadyl®, etodolac/Etogesic®, vedaprofen/Quadrisol-5®, tolfenamic acid/Tolfedine®). Yet others inhibit COX-2 exclusively (i.e., deracoxib/Deramax®, firocoxib/Previcox®, Robenacoxib/Onsior®, mavicoxib/Trocoxil®), while still another drug, acetaminophen, only weakly inhibits both

COX-1 and COX-2, but is able to inhibit COX-3 activity preferentially (Botting 2000).

Several in vitro studies investigating NSAID selective inhibition of the COX-1 and COX-2 isoenzymes have been published, but these are very difficult to interpret due to inconsistencies in the assays used (Kay-Mugford et al. 2000). Clinically, this information is confusing as it does not consider the pharmacokinetics of particular drugs and their concentrations in various tissues (Bertolini et al. 2001). Most NSAIDs that inhibit COX have been shown to result in diversion of arachidonate to the 5-LOX pathway. 5-LOX is principally found in polymorphonuclear cells, mast cells, monocytes, basophils, and B lymphocytes that are recruited during inflammatory and immune reactions (Bertolini et al. 2001). This enzyme catalyzes the initial step in leukotriene biosynthesis, which subsequently produces various eicosanoids, with LTB<sub>4</sub> being the most notable potent mediator of inflammation. The excessive production of leukotrienes has been implicated in the

creation of NSAID-induced ulcers (Hudson et al. 1993; Rainsford 1992). As always, however, the biological system is not clear-cut. While the LOX pathway is pro-inflammatory, there is also an anti-inflammatory pathway, which is discussed in more detail later (Serhan & Chiang 2004).

The contribution of the leukotrienes to the inflammatory process would seem to suggest that inhibition of both the COX and 5-LOX pathways by a therapeutic agent would result in an enhanced safety profile and may confer even greater analgesic efficacy due to broader anti-inflammatory and antinociceptive effects (Kirchner et al. 1997a). Data available show that dual-acting compounds are effective in arthritic models, where they also retain antithrombotic activity, produce little or no gastrointestinal damage, and do not adversely affect the asthmatic state (Bertolini et al. 2001).

A dual COX/5-LOX inhibitor called tepoxalin/Zubrin® underwent clinical trials and was approved for veterinary use in dogs (Kirchner et al. 1997b; Argentieri et al. 1994). Tepoxalin demonstrated gastrointestinal anti-inflammatory activity in mice, which supported the theory that 5-LOX inhibition may play a vital role in the prevention of NSAID-induced gastric inflammation (Kirchner et al. 1997b). Despite its approval in the United States, the manufacturer has voluntarily withdrawn the product from the market and it is no longer available.

### Prostaglandin Receptor Antagonism

Instead of limiting prostaglandin production through COX inhibition, a novel and more targeted therapeutic approach involves a class of drugs known as piprants, which are prostaglandin receptor antagonists (PRAs). Prostaglandin E<sub>2</sub> has been identified as a key mediator of the classic signs of inflammation including swelling, erythema, and pain through PGE<sub>2</sub>-mediated sensitization of peripheral neurons, as well as vasodilation and increased vascular permeability (Ricciotti & FitzGerald 2011). While PGE<sub>2</sub> exerts its effects via four different G protein-coupled receptors (EP1, EP2, EP3, and EP4), the EP4 receptor has been shown to play a dominant role in inflammation and pain, and may mediate central sensitization in chronic pain states. A specific EP4 receptor antagonist, grapiprant/Galliprant®, has been demonstrated to provide relief from arthritic pain in canine clinical patients and appears to act without affecting the synthesis and widespread activity of PGE<sub>2</sub> and other prostanoids (Kirkby Shaw et al. 2015).

### Analgesic and Anti-inflammatory Effects of NSAIDs

There is overwhelming evidence indicating that decreased PG synthesis due to COX inhibition and/or antagonism of PG activity at the EP4 receptor mediates NSAID-induced

analgesic and anti-inflammatory effects. Traditionally, NSAIDs have been considered peripheral-acting drugs that penetrate inflamed tissues, decrease PG-induced vasodilation and extravasation of protein exudates, and attenuate nociceptor sensitization. These local effects make them excellent analgesic choices for injuries with associated inflammation, as well as conditions such as synovitis, arthritis, cystitis, dermatitis, etc. (Chopra et al. 2000).

In addition to their peripheral effects, central analgesic effects have also been documented in several animal models of pain where NSAIDs have been shown to disrupt prostaglandin-mediated effects at both spinal and supraspinal levels (Chopra et al. 2000; Malmberg & Yaksh 1992; McCormack 1994; Yaksh et al. 1998). This central effect may account for the overall well-being and improved appetite that is often observed in patients receiving parenterally administered NSAIDs for relief of acute pain.

The relationship between COX inhibition and clinical relief of pain and inflammation in the intact animal is complex. Nociceptive pathways may involve either the COX-1 or COX-2 gene, and these genes are expressed in different locations and under different circumstances.

### NSAIDs and Fever

Just as the relationship between pain and the various activities of the COX system is complex, so too is the association between fever and the COX isoenzymes. The mechanisms leading to the generation of fever vary depending on the inciting factor, which may be peripheral (i.e., endotoxin) or central (i.e., endogenous pyrogens, such as IL-1). Interspecies variation is also substantial and the definitive role of the COXs in pyrexia remains to be clearly elucidated. Evidence suggests that COX-2 plays a role in endotoxin pyrexia while, based on the antipyretic effects of acetaminophen and aspirin, COX-1 and COX-3 appear to function in endogenous pyrexia (Botting 2000, 2003; Chandrasekharan et al. 2002).

### Other Pharmacologic Considerations for NSAID Use

Because of their high protein binding NSAIDs can displace other drugs from their plasma protein binding sites and potentially increase their plasma concentration. This is rarely a concern unless administered to animals with organ dysfunction or in those receiving other highly protein-bound medications with a narrow therapeutic index. Interference with the metabolism and excretion of certain coadministered drugs may occur, therefore verifying the safety of combination therapy is always mandatory.

Some NSAIDs may induce the syndrome of inappropriate secretion of ADH. Renal water reabsorption depends

on the action of ADH mediated by cyclic adenosine monophosphate (cAMP). As PGs exert a controlled negative feedback action on cAMP production, inhibition of PG synthesis results in above normal levels of cAMP with potential for enhanced ADH activity. In addition, the administration of a COX-2 selective NSAID may enhance sodium and water reabsorption. Clinically, both mechanisms may result in high specific gravity urine with dilutional hyponatremia. Urine volume may be decreased through this mechanism but without renal injury (Petersson et al. 1987; Dunn & Buckley 1986).

### Contraindications for NSAIDs

Nonsteroidal anti-inflammatory analgesics should not be administered to animals with acute renal insufficiency, hepatic insufficiency, dehydration, hypotension, conditions associated with low effective circulating volume (i.e., congestive heart failure, ascites), coagulopathies (i.e., factor deficiencies, thrombocytopenia, von Willebrand's disease), or evidence of gastric ulceration (i.e., vomiting with or without the presence of "coffee ground material", melena). Administration of NSAIDs following gastrointestinal surgery must be determined by the health of this organ at the time of surgery. As the COX-2 isoenzyme is important for healing, intuitively NSAIDs would be contraindicated where a compromised bowel is noted. Concurrent use of other NSAIDs (i.e., aspirin) or corticosteroids is not recommended. The COX-1 preferential NSAIDs are contraindicated in animals with spinal injury (including herniated intervertebral disc) due to the potential for hemorrhage and neurologic deterioration, and due to excessive bleeding at the surgical site should surgical treatment be pursued. The NSAIDs should never be administered to animals in shock, trauma cases at the time of presentation, or animals with evidence of hemorrhage (i.e., epistaxis, hemangiosarcoma, head trauma). Animals with severe or poorly controlled asthma, or other types of moderate to severe pulmonary disease, may deteriorate with NSAID administration. Aspirin has been documented to exacerbate asthma in human patients, but COX-2-specific NSAIDs did not result in worsening of clinical signs (West & Fernandez 2003). It is not known whether animals may be affected in this way. Although administration of NSAIDs in head trauma, pulmonary diseases, or thrombocytopenia is generally contraindicated, COX-2 preferential NSAIDs (i.e., meloxicam, etodolac, carprofen, tolafenamic acid, and deracoxib) may prove to be safe with further study. Due to inhibition of PG activity, the NSAIDs may be detrimental to reproductive function. Indomethacin may block PG activity in pregnant women, resulting in cessation of labour, premature closure of the ductus arteriosus in the

fetus, and disruption of fetal circulation (DuBois et al. 1998). These effects may occur in animals, therefore NSAIDs should not be administered during pregnancy. As COX-2 induction is necessary for ovulation and subsequent implantation of the embryo (DuBois et al. 1998), NSAIDs should also be avoided in breeding females during this stage of the reproductive cycle. As previously mentioned, the COX-2 isoenzyme is required for maturation of the embryological kidney so administration to lactating mothers should be avoided.

### NSAIDs Used in Veterinary Medicine

There are few data regarding clinical pharmacology of NSAIDs in exotic species, thus care must be taken when extrapolating from common domestic species.

#### Meloxicam

Meloxicam is a COX-2 preferential NSAID approved for use in dogs in Australasia, Europe, and North America. The parenteral formulation is approved for cats in the United States and Australasia. Its use in cats in Canada is under investigation, with completed studies indicating safety and efficacy. Its use in horses is also under investigation, with pharmacokinetic studies indicating that the half-life is shorter and clearance greater than in the dog, suggesting that dosing more than once a day may be necessary (Sinclair et al. 2003).

Studies indicate no renal or hepatic abnormalities with acute administration and minimal to no antithrombotic activity, suggesting hemostasis in normal animals may not be a problem (Mathews et al. 1999; Poulsen & Justus 1999). Few adverse reactions have been documented and most involve the gastrointestinal tract. A recent study showed no difference in gastric erosions over saline placebo when meloxicam was administered at 0.1 mg/kg for 3 days post electrical stimulation (i.e., surgical simulation) under anesthesia. However, corticosteroids plus meloxicam in this study resulted in significant gastric erosions (Boston et al. 2003). A case report of combination aspirin and meloxicam in a dog resulted in duodenal perforation (Reed 2002). This case illustrates the importance of COX-2 in intestinal protection when aspirin is coadministered, and reinforces that different NSAIDs should not be administered concurrently. Analgesia is excellent when meloxicam is combined with an opioid.

In recent years, meloxicam has become the most widely used NSAID in exotic and zoo animals, including reptiles, birds, and numerous mammalian species. Drug use remains off-label, with doses largely extrapolated from domestic animal pharmacokinetic and pharmacodynamic studies. Species-specific studies are beginning to appear in

the literature and should be consulted wherever possible prior to prescribing meloxicam or any NSAID.

### **Carprofen**

Although classified as a NSAID, carprofen administration to beagle dogs did not inhibit PGE<sub>2</sub>, 12-hydroxyeicosatetraenoic acid, or thromboxane B<sub>2</sub> synthesis in an experimental study utilizing subcutaneous tissue cage fluids (McKellar et al. 1994b). It was concluded that the principle mode of action of carprofen must be by mechanisms other than cyclooxygenase or 12-lipoxygenase inhibition. However, more recent studies indicate that it is a COX-2 preferential NSAID (Kay-Mugford et al. 2000; Ricketts et al. 1998). Carprofen is approved for perioperative and chronic pain management in dogs in Australasia, Europe, and North America. Carprofen is approved for single dose, perioperative use in cats in Europe, and is licensed for use in horses in the United Kingdom. In sheep, carprofen (0.7 mg/kg IV) resulted in plasma concentrations of 1.5 µg/mL, similar to those required to confer analgesia in horses, for up to 48 hours (Welsh et al. 1992). However, analgesia was not assessed in this sheep study (Welsh et al. 1992). Antithromboxane activity is minimal, suggesting that induced coagulopathy may not be a problem in animals with intact hemostatic mechanisms (Poulsen & Justus 1999; McKellar et al. 1990).

### **Ketoprofen**

Ketoprofen is approved for postoperative and chronic pain in both dogs and cats in Europe and Canada. Ketoprofen is also approved for use in horses and ruminants. As ketoprofen is an inhibitor of both COX-1 and COX-2, adverse effects are a potential problem requiring careful patient selection. Although several studies using ketoprofen preoperatively indicate its effectiveness in controlling postoperative pain, a general consensus among veterinarians has restricted its use primarily to the postoperative period to reduce the potential for hemorrhage (Mathews et al. 1999; Lobetti & Joubert 2000; Pibarot et al. 1997). Ketoprofen should not be administered to patients with risk factors for hemorrhage. It is often administered to animals immediately after orthopedic procedures (i.e., fracture repair, cruciate repair, onychectomy).

### **Deracoxib**

Deracoxib is a COX-2 specific inhibitor. It is approved in the United States and Canada for control of postoperative pain and inflammation associated with orthopedic surgery in dogs. The incidence of vomiting and diarrhea were similar in dogs receiving deracoxib compared to dogs receiving placebo in a perioperative field trial, and overall the drug was well tolerated and effective (Novartis Animal Health

USA 2004). It was also shown to be effective in attenuating lameness in dogs with urate crystal-induced synovitis after prophylactic and therapeutic administration (McCann et al. 2004; Millis et al. 2002). This group of NSAIDs appeared to be gastroprotective in human patients when compared to the less COX-2 specific NSAIDs, when used for 8 days to 3 months (Silverstein et al. 2000). However, more recent studies in humans indicate these NSAIDs cannot guarantee gastroprotection with chronic use. Furthermore, in a canine study comparing the gastrointestinal safety profile of licofelone (a dual inhibitor) to rofecoxib (another specific COX-2 inhibitor), rofecoxib was found to induce significant gastric and gastroduodenal lesions (Moreau et al. 2005).

### **Diclofenac**

Diclofenac is available worldwide as several different human and veterinary formulations. It is a useful anti-inflammatory and analgesic drug, and has been studied for antimicrobial activity (Dutta et al. 2007). Diclofenac has been observed to cause severe hepatic and nephrotoxicity in many species of birds and its use, or accidental ingestion from carcasses, should be avoided (Jayakumar et al. 2010; Jain et al. 2009; Naidoo et al. 2009; Hussain et al. 2008; Taggart et al. 2007; Oaks et al. 2004).

### **Firocoxib**

Firocoxib is available as an oral paste and injectable for horses and as an oral formulation for dogs. The efficacy and adverse events appear similar to other coxib-class NSAIDs (US Food and Drug Administration 2011). There are few reports of firocoxib use in birds, reptiles, or exotic mammals in the literature.

### **Robenacoxib**

Robenacoxib is available as an oral and injectable formulation for treatment of pain and inflammation in both dogs and cats. Specific label indications and doses vary by country. The efficacy and adverse events appear similar to other coxib-class NSAIDs.

### **Mavicoxib**

Mavicoxib is a long-acting NSAID approved outside the United States for use in dogs. The dosing interval is usually 2 weeks between doses 1 and 2, then 4 weeks between subsequent doses. The elimination half-life in healthy dogs is approximately 2 weeks (range 7.9–38.8 days) (Cox et al. 2010). Significant individual and breed-associated differences have been observed but effects on drug safety and effectiveness have yet to be determined (Cox et al. 2011).

### Tolfenamic Acid

Tolfenamic acid is approved for use in cats and dogs in Europe and Canada for controlling acute postoperative and chronic pain. The dosing schedule is 3 days on and 4 days off, which must be strictly adhered to. Reported adverse effects are diarrhea and occasional vomiting. Tolfenamic acid has significant anti-inflammatory and antithromboxane activity, therefore post-traumatic and surgical hemostasis may be compromised during active bleeding after administration of this NSAID (McKellar et al. 1994b).

### Flunixin Meglumine

Flunixin meglumine is a COX-1 and COX-2 inhibitor that is approved for use in dogs in Europe but not North America. It is also approved for use in ruminants and horses, and is commonly used for equine colic pain.

### Phenylbutazone

Phenylbutazone is approved for use in horses, cattle, and dogs in North America. Since safer NSAIDs are approved for dogs, phenylbutazone is not recommended for this species. In horses, there is high risk of gastric ulceration and nephrotoxicity, where signs of toxicity may progress from inappetence and depression to colic, gastrointestinal ulceration, and weight loss (MacAllister et al. 1993; Collins & Tyler 1984; Snow et al. 1981). Phenylbutazone has a prolonged elimination half-life in cattle, ranging from 30 to 82 hours (DeBacker et al. 1980; Arifah & Lees 2002).

### Aspirin

Aspirin is primarily a COX-1 inhibitor. It has been most commonly used as an analgesic for osteoarthritic pain in dogs. It is also available in proprietary combinations with various opioids (aspirin plus codeine or aspirin plus oxycodone) to achieve a synergistic effect for the treatment of moderate pain. It is also used as an antipyretic and anticoagulant in dogs and cats. Aspirin has also been recommended in cattle (Gingerich et al. 1975).

### Grapiprant

Grapiprant is a non-COX-inhibiting NSAID that acts as an EP4 PG receptor antagonist downstream of traditional NSAIDs in the arachidonic acid cascade. It is approved for treatment of pain and inflammation associated with osteoarthritis in dogs and is supplied in 20-, 60-, and 100-mg flavored tablets. In a clinical trial submitted as part of the regulatory approval process in the United States, grapiprant was found to be effective compared to placebo, although there was no positive control. In the same study, grapiprant was generally well tolerated by dogs, with intermittent diarrhea, vomiting, and anorexia the most commonly reported adverse effects (Rausch-Derra et al. 2016).

More studies are needed to demonstrate grapiprant's analgesic efficacy in osteoarthritis and other types of pain and inflammation in dogs and to compare its safety to other approved COX-2 inhibiting NSAIDs.

## NSAIDs Not Approved for Use in Veterinary Medicine (Off-label Use)

### Ketorolac

Ketorolac is a COX-1 and COX-2 inhibitor that is included for the benefit of those working in the research setting associated with human hospitals where the availability of ketorolac is more likely than other NSAIDs. Adverse gastrointestinal effects are common.

### Acetaminophen

Acetaminophen is a COX-3 inhibitor with minimal COX-1 and COX-2 effects. It should not be administered to feline species due to deficient glucuronidation of acetaminophen in these species (Court & Greenblatt 1997). Reptiles are known to be particularly sensitive to the toxic effects of acetaminophen and this has been exploited to combat the invasive brown treesnake on the island of Guam by lacing dead mice with 80 mg of acetaminophen and baiting snake habitats. A recent study has confirmed that the standard 80-mg dose used operationally to kill brown treesnakes produced acute methemoglobinemia, with severe hypoxia and respiratory failure cited as the probable cause of death (Mathies & Mauldin 2020). Based on these findings, acetaminophen should not be administered to any reptile at any dose.

### Dipyrone

Dipyrone is a COX-3 inhibitor and is approved for use in cats and dogs in Europe and Canada. Dipyrone should be given intravenously to avoid the irritation experienced when given intramuscularly. The analgesia produced is not usually adequate for moderate to severe postoperative pain, and dipyrone is reserved for use as an antipyretic in cases where other NSAIDs are contraindicated. Dipyrone induces blood dyscrasias in humans, but this has not been reported in animals.

## Alpha-2 Adrenergic Receptor Agonists and Antagonists

### Introduction

The use of alpha-2 adrenergic agonists in veterinary medicine began following the synthesis of xylazine in 1962. Early reports of the sedative and anesthetic sparing qualities of xylazine predated the elucidation of its mechanism

of action in 1981. Alpha-2 adrenoreceptors have been identified in the CNS, cardiovascular, respiratory, renal, endocrine, gastrointestinal, and hemotologic systems, resulting in widespread drug effects (Aantaa et al. 1995). Most FDA-CVM approved alpha-2 agonists carry label indications as sedatives and analgesics.

Alpha-2 adrenoceptors are linked to Gi-protein second messengers (Aantaa et al. 1995). These are similar to those used by many opioid receptor subtypes and in fact opioid and alpha-2 agonists usually have additive or synergistic effects (Maze & Tranquilli 1991). Alpha-2 adrenoreceptors are classically described as being located presynaptically at noradrenergic neurons, exerting an inhibitory feed-back role on the release of subsequent norepinephrine (Maze & Tranquilli 1991). This results in decreased sympathetic nervous system efferent activity and is probably related to the decreased vigilance, decreased anesthetic requirements, and decreased heart rate and blood pressure observed following administration of these drugs to most species. Alpha-2 adrenoreceptors are also found in the vascular smooth muscle (a non-presynaptic site) and when activated result in vasoconstriction. This results in increased vascular resistance and will result in increased baroreceptor-mediated vagal tone. The result is slowing of heart rate and decreasing cardiac output, but blood pressure remains within physiologic normal values. Confusion often exists about the clinical effect of alpha-2 agonist administration (e.g., hypertension [post-synaptic] vs. hypotension [pre-synaptic]) expected in a patient. The net clinical result will vary with route of administration, dose, species, and the duration of time following the administration.

Alpha-2 agonist doses vary at least 10-fold across species. Pigs tend to have the highest requirements, followed by cats, dogs, horses, and finally ruminants. Breed and sex differences also exist within cattle that should be appreciated.

## Alpha-2 Adrenergic Agonist Effects

### Central Nervous System

Alpha-2 adrenergic agonists exert many of their inhibitory effects on CNS function through inhibiting norepinephrine release from sympathetic neurons (Maze & Tranquilli 1991). Inhibition of intraneuronal transmission is also responsible for muscle relaxation observed following alpha-2 adrenergic agonist administration. Analgesia is mediated by spinal and supraspinal alpha-2 adrenergic receptors. Agonist binding modulates afferent activity at a spinal level and increases the diffuse noxious inhibitory control system activity. The net result is sedation, reduced anesthetic requirements, reduced stress responses, and analgesia. It should be noted that alpha-2 agonists, like all sedatives and tranquilizers, are not

anesthetics. Although animals can appear in a sleep-like state, they may become aroused by noxious stimulation and may become defensive. Additionally, on rare occasions, paradoxical behavior (aggression rather than sedation) may be noted. Accidental intracarotid injection of alpha-2 adrenergic agonists will induce seizure-like activity and must be avoided.

### Emetic Center

Alpha-2 agonists are predictable emetics in cats and dogs, especially at high doses. This is due to activation of the chemoreceptor trigger zone of the area postrema (Hikasa et al. 1992). Dopaminergic blocking agents do not prevent alpha-2 induced emesis.

### Thermoregulation

Alpha-2 agonists will often cause changes in thermoregulation. The effect is usually a decrease; however, increases can be seen when animals are placed in warm environments. This is especially a concern when alpha-2 agonists are used for capture of cattle or other hoof-stock. If possible, body temperature should be monitored for 12–24 hours following sedation or administration of an alpha-2 adrenergic antagonist should be considered. In smaller animals, hypothermia is more common and may be due to decreased metabolic activity accompanying sedation as well as decreased thermoregulatory control. It is usually not a life-threatening problem if managed appropriately.

### Eyes

Alpha-2 agonists generally cause mild miosis to little change in pupil diameter and a mild decrease in intraocular pressure (IOP) (Verbruggen et al. 2000). This class of drugs is relatively contraindicated, with increases in IOP or corneal lacerations because of the probability of inducing vomiting (which causes further increases in IOP) in those species that can vomit.

### Respiratory System

Most alpha-2 adrenergic receptor agonists can cause decreased responsiveness to CO<sub>2</sub>, especially in higher doses. This effect is compounded by the coadministration of opioids or anesthetic agents and is of concern during anesthesia. In some species alpha-2 agonists do not normally depress respiration to the point of creating hypoxia and cyanosis. In fact, arterial blood gas values for PaO<sub>2</sub> are usually normal. The bluish color of the mucous membranes that may be noticed on occasion is usually due to slowed venous blood flow accompanying vasoconstriction. As the capillary transit time increase, oxygen extraction increases and notable hemoglobin desaturation will appear as a blue(ish) color. Oxygen therapy will often resolve this

condition. However, in sheep and possibly some other species, xylazine administration has been shown to result in central hypoxemia related to pulmonary changes (Celly et al. 1997, 1999).

### Cardiovascular System

Intravenous administration of alpha-2 adrenergic agonists typically results in rapid and pronounced vasoconstriction (Pypendop & Verstegen 1998). If heart rate and cardiac output do not decrease (as can be seen with atropine administration), blood pressure would increase impressively (Alibhai et al. 1996; Short 1991). However, normal baroreceptor reflexes attempt to maintain blood pressure within physiologic limits by increasing vagal tone and slowing heart rate. Heart rate may decrease by 50–75% in some individuals. The cardiac rhythm is often a slow sinus rhythm characterized by two or three sequential beats followed by a long sinus pause. Occasionally, second degree atrioventricular blockade will be seen. Both rhythms are often responsive to antimuscarinic administration (block the vagal reflex) and high heart rates (and blood pressures) will result. Intramuscular or oral administration tends to decrease the incidence of profound bradycardia.

After the initial direct vasoconstriction occurs, vasodilation and reduction in blood pressure may be seen. This is more prominent in primate species. The alpha-2 adrenergic receptor agonists decrease sympathetic nervous system efferent activity, which results in decreased vasomotor tone and heart rate. Alpha-2 adrenergic agonists have historically been used in humans as antihypertensive agents because they reduce vasomotor tone and block the reflex increase in heart rate that can accompany alpha-1 antagonists.

Most of the contraindications to alpha-2 adrenergic receptor agonist administration are related to their cardiovascular side effects. As a general rule, this class of drugs should not be administered to animals that do not have normal healthy cardiovascular systems and/or are exercise intolerant. There are some exceptions, but a thorough understanding of the underlying disease is required and appropriate monitoring is necessary.

### Urinary Tract

Alpha-2 adrenergic receptor agonists increase urine output by increasing production of dilute urine (Saleh et al. 2005; Grimm et al. 2001). This is primarily related to inhibition of ADH release and/or synthesis as well as changes in renal hemodynamics (Saleh et al. 2005).

### Gastrointestinal Tract

Animals may have complications associated with decreased propulsive activity, including colic and bloat, although this is unusual in most species (Thompson et al. 1991).

Longer-acting drugs are more likely to result in a problem. Patients should be observed following alpha-2 adrenergic receptor agonist administration for signs of abdominal distension.

### Endocrine

A classic neuroendocrine response is hyperglycemia following alpha-2 agonist administration (Abdel el Motal & Sharp 1985; Osman & Nicholson 1991). This is due to a decrease in insulin release. It is usually transient and not clinically significant, although alpha-2 agonists should not be used to sedate animals for glucose curves.

Since alpha-2 agonists reduce sympathetic activity and inhibit the stress response, neuroendocrine markers of the stress response should be affected. Cortisol levels are usually decreased following alpha-2 agonist administration and may not be reliable as indicators of stress or pain (Brearley et al. 1990; Sanhoury et al. 1992).

### Specific Drugs

Several alpha-2 adrenergic receptor agonists are approved for veterinary use. Generally, their pharmacologic actions will be similar, but the duration of action and species compatibility will vary. All alpha-2 agonists are potent and potentially dangerous following accidental human ingestion or injection. Care should be taken when handling syringes loaded with these drugs and medical help should be sought immediately if exposure occurs.

#### Xylazine

Xylazine is the prototypical veterinary alpha-2 adrenergic receptor agonist. Another alpha-2 agonist, clonidine, has been used in humans as an antihypertensive agent and is often used as the prototypical drug in research applications. Xylazine has been administered to many different species, both domestic and exotic. It is readily available and relatively inexpensive.

Xylazine has a shorter duration of action than many of the other drugs. Typical doses will result in muscle relaxation and sedation of horses for 45–60 minutes. This can be advantageous when performing field anesthesia/sedation when a rapid recovery is desired. Xylazine is often combined with opioids such as butorphanol to enhance sedative and analgesic qualities.

#### Detomidine

Detomidine is a longer-acting alpha-2 agonist approved for use in horses as a sedative. It is commonly administered when profound, long-lasting sedation is needed. When used in high doses as a preanesthetic, low respiratory rates may accompany induction. Detomidine is not used in small

animals but there are a number of reports evaluating its use in combination with other agents in large exotic species, including rhinoceros and zebras.

### **Medetomidine**

Medetomidine is formulated as a racemic mixture that was originally approved as a sedative/analgesic in dogs. Though it is no longer widely available except as formulations prepared for wildlife capture and restraint, it has been extensively evaluated in all domestic species and many exotic and zoological species. It is extremely selective for the alpha-2 receptor and binds it avidly. Atipamezole was developed as the specific antagonist for medetomidine for this reason.

### **Romifidine**

Romifidine is an alpha-2 adrenergic receptor agonist that is approved for use in horses. It has been evaluated in other species and appears to be relatively safe but offers few advantages over other approved products. Some equine clinicians believe it is a good sedative without causing excessive ataxia.

### **Dexmedetomidine**

Dexmedetomidine contains only the pure dextrorotatory enantiomer of racemic medetomidine, which appears to be responsible for all of the clinically relevant properties of the drug. Due to the absence of the inactive levorotatory enantiomer, dexmedetomidine is twice as potent as medetomidine. It is currently marketed as a solution with half the strength of medetomidine, meaning that equivalent doses of dexmedetomidine and medetomidine have the same volume. The sedative, analgesic, and anesthetic-sparing effects of both drugs appear to be similar when administered at equivalent doses in dogs and cats (Ansah et al. 1998; Granholm et al. 2006, 2007; Kuusela et al. 2000). Hemodynamic side effects and reversal characteristics with administration of the specific antagonist atipamezole also appear similar. Dexmedetomidine has been evaluated in combination with a variety of other injectable agents, including benzodiazepines, dissociatives, opioids, and alfaxalone, in many zoo, exotic, and wildlife species.

## **Alpha-2 Adrenergic Receptor Antagonists**

One big advantage alpha-2 adrenoreceptor agonists have over other sedative/tranquilizers such as acepromazine is reversibility. It is possible to administer an antagonist and within minutes animals will regain function and be able to be released. Antagonist administration will usually result in opposite hemodynamic effects (e.g., acute vasodilation and tachycardia) and they should not be administered in a

cavalier fashion. Many of the difficulties (including some deaths) occur with reversal. Also, alpha-2 agonist mediated analgesia is reversed, necessitating administration of other analgesic classes to painful animals. Some species (e.g., domestic horses) are not routinely reversed because of the potential for uncontrollable activity and possible self-trauma.

### **Yohimbine**

Yohimbine is a plant-derived compound that has been used by humans for centuries because of its performance enhancing properties. It is relatively effective for reversing xylazine in horses, dogs, and cats, but is less effective in ruminants. Additionally, yohimbine has some stimulant actions that may result in excitation on recovery. Yohimbine has not been effective at reversing newer agents, such as medetomidine.

### **Tolazoline**

Tolazoline was originally used as a therapy for human infants suffering from pulmonary hypertension. It is a relatively nonselective alpha-receptor blocker and is useful for reducing pulmonary vascular resistance. It is also effective at antagonizing xylazine sedation. It is most commonly used for equine and ruminant species. It appears more effective in ruminant species than yohimbine.

### **Atipamezole**

Atipamezole is the marketed antagonist for medetomidine and dexmedetomidine. It is effective at reversing all available alpha-2 adrenoreceptor agonists and its use is limited only by cost. It is relatively selective for alpha-2 adrenergic receptors and usually does not cause excessive stimulation, although excitation may occur. Atipamezole is approved for use in dogs, but it is used in other species commonly. Under most circumstances, it is best given by the intramuscular route except in emergency situations.

## **Phenothiazines and Butyrophenones**

### **Introduction**

The phenothiazines and butyrophenones produce an array of behavioral, autonomic, and endocrine effects and have been used clinically in numerous domestic and wild species. Their tranquilizing effects are mediated by antagonism of dopamine receptors (primarily the D2 subtype) located in the cerebral cortex, basal ganglia, and limbic system. In addition, other antidopamine effects are noted in the hypothalamus (increased prolactin secretion and impaired thermoregulation), brainstem (impaired vasomotor reflexes), and chemoreceptor trigger zone of the

medulla (antiemesis). Both drug classes also have varying antagonistic effects at adrenergic (alpha-1 and alpha-2), serotonergic (5-hydroxytryptamine receptors), muscarinic (M1), and histaminergic (H1) receptor systems. It is important to note that they do not possess any inherent analgesic properties and their effects are not reversible.

### **Acepromazine**

Acepromazine is the most widely used phenothiazine tranquilizer in veterinary medicine. In dogs and cats it is commonly combined with an opioid to produce a state historically referred to as “neuroleptanalgesia”. Such combinations are suitable to provide chemical restraint for short, noninvasive procedures or as preanesthetic medication prior to induction of general anesthesia. Acepromazine administration in the preanesthetic period will cause dose-dependent reductions in both injectable and inhalant anesthetic requirements in these species. In horses and cattle, the drug is used primarily for its anti-anxiety effects. In susceptible pigs, acepromazine has been shown to prevent or reduce the onset of halothane-induced malignant hyperthermia. In certain wild and exotic species, the combination of acepromazine and the potent opioid etorphine (Large Animal Immobilon) has been used for immobilization and anesthesia.

The cardiovascular effects of acepromazine include decreases in mean arterial blood pressure in the range of 20–30% accompanied by dose-dependent reductions in stroke volume and cardiac output. Isoflurane, due to its potent vasodilatory effects, appears to potentiate acepromazine-induced hypotension. At clinically relevant doses, heart rate may not change appreciably or may increase slightly. At very high doses, bradycardia and sinoatrial block may occur. Acepromazine has also been shown to reduce the arrhythmogenic effects of epinephrine and halothane.

In general, effects on pulmonary function (i.e., oxygenation and ventilation) tend to be minimal in conscious animals, although respiratory rate may decrease somewhat. Acepromazine has been shown to dose-dependently decrease hematocrit by as much as 20–50% in dogs and horses due to splenic sequestration. This effect occurs within 30 minutes and appears to persist for at least 2 hours. The drug also decreases platelet aggregation but the clinical hemostatic significance of this appears minimal.

Regarding the gastrointestinal system, acepromazine has been shown to have antiemetic effects when administered 15 minutes prior to morphine, hydromorphone, or oxymorphone. Lower esophageal sphincter tone is reduced, which may lead to an increased risk of gastric reflux, though this has not been proven. Decreased gastrointestinal motility and delayed gastric emptying have been demonstrated in horses.

Renal blood flow and glomerular filtration rate appear to be well-maintained in dogs receiving acepromazine and isoflurane. In cats under halothane anesthesia, urethral pressure has been shown to decrease by 20% with acepromazine administration. There are anecdotal reports of penile prolapse/priapism in stallions. The magnitude and duration of the protrusion appear to be dose-dependent and this side effect appears to be mediated by alpha-adrenergic antagonism. Also, because of its antihistaminergic effects, acepromazine is not suitable for allergic skin-testing.

### **Azaperone**

Azaperone is classified as a butyrophenone and its only approved indication is for control of aggression associated with mixing or re-grouping of swine. It is also used as a preanesthetic agent in pigs and has clinical properties and side effects similar to acepromazine. In various wild and exotic species azaperone is used in combination with potent opioids such as etorphine or carfentanil to produce immobilization or anesthesia.

## **References**

- Aantaa R, Marjamaki A, Scheinin M. 1995. Molecular pharmacology of alpha 2-adrenoceptor subtypes. *Annals of Medicine* 27(4):439–449.
- Abdel el Motal SM, Sharp GW. 1985. Inhibition of glucose-induced insulin release by xylazine. *Endocrinology* 116(6):2337–2340.
- Algotsson L, Messeter K, Rosen I, et al. 1992. Effects of nitrous oxide on cerebral haemodynamics and metabolism during isoflurane anaesthesia in man. *Acta Anaesthesiologica Scandinavica* 36(1):46–52.
- Alibhai HI, Clarke KW, Lee YH, et al. 1996. Cardiopulmonary effects of combinations of medetomidine hydrochloride and atropine sulphate in dogs. *The Veterinary Record* 138(1):11–13.
- Ambros B, Duke-Novakovski T, Pasloske KS. 2008. Comparison of the anesthetic efficacy and cardiopulmonary effects of continuous rate infusions of alfaxalone-2-hydroxypropyl-beta-cyclodextrin and propofol in dogs. *American Journal of Veterinary Research* 69:1391–1398.

- American Veterinary Medical Association Guidelines for the Euthanasia of Animals, 2020 Edition.
- Ansah OB, Raekallio M, Vainio O. 1998. Comparison of three doses of dexmedetomidine with medetomidine in cats following intramuscular administration. *Journal of Veterinary Pharmacology and Therapeutics* 21:380–387.
- Argentieri DC, Ritchie DM, Ferro MP, et al. 1994. Tepoxalin: a dual cyclooxygenase/5-lipoxygenase inhibitor of arachidonic acid metabolism with potent anti-inflammatory activity and a favorable gastrointestinal profile. *The Journal of Pharmacology and Experimental Therapeutics* 271:1399–1408.
- Arifah AK, Lees P. 2002. Pharmacodynamics and pharmacokinetics of phenylbutazone in calves. *Journal of Veterinary Pharmacology and Therapeutics* 25:299–309.
- Barnhart MD, Hubbell JAE, Muir WW, et al. 2000. Pharmacokinetics, pharmacodynamics, and analgesic effects of morphine after rectal, intramuscular, and intravenous administration in dogs. *American Journal of Veterinary Research* 61:24–28.
- Barter LS, Ilkiw JE, Pypendop BH, et al. 2004. Evaluation of the induction and recovery characteristics of anesthesia with desflurane in cats. *American Journal of Veterinary Research* 65(6):748–751.
- Bednarski RM, Gaynor JS, Muir WW 3rd. 1993. Vaporizer in circle for delivery of isoflurane to dogs. *Journal of the American Veterinary Medical Association* 202(6):943–948.
- Berry, SH. 2015. Injectable anesthetics. In: *Veterinary Anesthesia and Analgesia: The Fifth Edition of Lumb and Jones* (KA Grimm, KA Lamont, WJ Tranquilli, SA Greene, SA Robertson, eds.). Ames: Wiley Blackwell.
- Bertolini A, Ottani A, Sandrini M. 2001. Dual acting anti-inflammatory drugs: a reappraisal. *Pharmacological Research* 44:437–450.
- Bini G, Vettorato E, De Gennaro C, et al. 2018. A retrospective comparison of two analgesic strategies after uncomplicated tibial plateau levelling osteotomy in dogs. *Veterinary Anaesthesia and Analgesia* 45(4):557–565.
- Bitti FS, Campagnol D, Rangel JP, et al. 2017. Effects of three methadone doses combined with acepromazine on sedation and some cardiopulmonary variables in dogs. *Veterinary Anaesthesia and Analgesia* 44(2):237–245.
- Boidin MP. 1985. Serum levels of cortisol in man during etomidate, fentanyl and air anesthesia, compared with neurolept anesthesia. *Acta Anaesthesiologica Belgica* 36(2):79–87.
- Borland K, Garcia E. 2019. Case report: possible opioid-induced hyperalgesia in a cat. *Companion Anim* 24:246–248.
- Boston SE, Moens NM, Kruth SA, et al. 2003. Endoscopic evaluation of the gastroduodenal mucosa to determine the safety of short-term concurrent administration of meloxicam and dexamethasone in healthy dogs. *American Journal of Veterinary Research* 64:1369–1375.
- Botting RM. 2000. Mechanism of action of acetaminophen: is there a cyclooxygenase 3? *Clinical Infectious Diseases : An Official Publication of the Infectious Diseases Society of America* 31(Suppl 5):S202–210.
- Botting R. 2003. COX-1 and COX-3 inhibitors. *Thrombosis Research* 110:269–272.
- Bowdle TA. 1998. Adverse effects of opioid agonists and agonist-antagonists in anaesthesia. *Drug Safety* 19:173–189.
- Branson K. 2007. Injectable and alternative anesthetic techniques. In: *Lumb and Jones' Veterinary Anesthesia and Analgesia*, 4th ed. (WJ Tranquilli, JC Thurmon, KA Grimm, eds.). Ames: Blackwell Publishing.
- Branson KR, Gross ME, Booth NH. 2001. Opioid agonists and antagonists. In: *Veterinary Pharmacology and Therapeutics* (HR Adams, ed.), p. 274. Ames: Iowa State Press.
- Brearley JC, Dobson H, Jones RS. 1990. Investigations into the effect of two sedatives on the stress response in cattle. *Journal of Veterinary Pharmacology and Therapeutics* 13(4):367–377.
- Briggs SL, Sneed K, Sawyer DC. 1998. Antinociceptive effects of oxymorphone-butorphanol-acepromazine combination in cats. *Veterinary Surgery* 27:466–472.
- Brown EN, Lydic R, Schiff ND. 2010. General anesthesia, sleep and coma. *New England Journal of Medicine* 363:2638–2650.
- Carroll GL, Hooper RN, Boothe DM, et al. 1999. Pharmacokinetics of fentanyl after intravenous and transdermal administration in goats. *American Journal of Veterinary Research* 60:986–991.
- Caulkett NA, Cribb PH, Haigh JC. 2000. Comparative cardiopulmonary effects of carfentanil-xylazine and medetomidine-ketamine used for immobilization of mule deer and mule deer/white-tailed deer hybrids. *Canadian Journal of Veterinary Research – Revue Canadienne de Recherche Veterinaire* 64:64–68.
- Celly CS, McDonell WN, Young SS, et al. 1997. The comparative hypoxaemic effect of four alpha 2 adrenoceptor agonists (xylazine, romifidine, detomidine and medetomidine) in sheep. *Journal of Veterinary Pharmacology and Therapeutics* 20(6):464–471.
- Celly CS, Atwal OS, McDonell WN, et al. 1999. Histopathologic alterations induced in the lungs of sheep by use of alpha2-adrenergic receptor agonists. *American Journal of Veterinary Research* 60(2):154–161.
- Chandrasekharan NV, Dai H, Roos KL, et al. 2002. COX-3, a cyclooxygenase-1 variant inhibited by acetaminophen and other analgesic/antipyretic drugs: cloning, structure, and expression. *Proceedings of the National Academy of Sciences of the United States of America* 99:13926–13931.

- Chau PL. 2010. New insights into the molecular mechanisms of general anaesthetics. *British Journal of Pharmacology* 161(2):288–307.
- Chism JP, Rickert DE. 1996. The pharmacokinetics and extra-hepatic clearance of remifentanyl, a short acting opioid agonist, in male beagle dogs during constant rate infusions. *Drug Metabolism and Disposition* 24:34–40.
- Chopra B, Giblett S, Little JG, et al. 2000. Cyclooxygenase-1 is a marker for a subpopulation of putative nociceptive neurons in rat dorsal root ganglia. *European Journal of Neuroscience* 12:911–920.
- Christie MJ, Connor M, Vaughan CW, et al. 2000. Cellular actions of opioids and other analgesics: Implications for synergism in pain relief. *Clinical and Experimental Pharmacology and Physiology* 27:520–523.
- Clarke KW. 1999. Desflurane and sevoflurane. New volatile anesthetic agents. *The Veterinary Clinics of North America. Small Animal Practice* 29(3):793–810.
- Collins LG, Tyler DE. 1984. Phenylbutazone toxicosis in the horse: a clinical study. *Journal of the American Veterinary Medical Association* 184:699–703.
- Colvin LA, Bull F, Hales TG. 2019. Perioperative opioid analgesia – when is enough too much? A review of opioid-induced tolerance and hyperalgesia. *Lancet* 13;393(10180):1558–1568.
- Court MH, Greenblatt DJ. 1997. Molecular basis for deficient acetaminophen glucuronidation in cats: An interspecies comparison of enzyme kinetics in liver microsomes. *Biochemical Pharmacology* 53:1041–1047.
- Cox SR, Lesman SP, Boucher JF, et al. 2010. The pharmacokinetics of mavacoxib, a long-acting COX-2 inhibitor, in young adult laboratory dogs. *Journal of Veterinary Pharmacology and Therapeutics* 33(5):461–470.
- Cox SR, Liao S, Payne-Johnson M, et al. 2011. Population pharmacokinetics of mavacoxib in osteoarthritic dogs. *Journal of Veterinary Pharmacology and Therapeutics* 34(1):1–11.
- Daghfous R, el Aidli S, Sfaxi M, et al. 2003. Halothane-induced hepatitis. 8 case reports. *La Tunisie Medicale* 81(11):874–878.
- Davidson G. 2001. To benzoate or not to benzoate: Cats are the question. *International Journal of Pharmaceutical Compounding* 5(2):89–90.
- DeBacker P, Braeckman R, Belpaire F. 1980. Bioavailability and pharmacokinetics of phenylbutazone in the cow. *Journal of Veterinary Pharmacology and Therapeutics* 3:29–33.
- Deng M, Chen SR, Chen H, et al. 2019.  $\alpha 2\delta$ -1-bound N-methyl-D-aspartate receptors mediate morphine-induced hyperalgesia and analgesic tolerance by potentiating glutamatergic input in rodents. *Anesthesiology* 130(5):804–819.
- Dodam JR, Kruse-Elliott KT, Aucoin DP, et al. 1990. Duration of etomidate-induced adrenocortical suppression during surgery in dogs. *American Journal of Veterinary Research* 51(5):786–788.
- Dodam JR, Cohn LA, Durham HE, et al. 2004. Cardiopulmonary effects of medetomidine, oxymorphone, or butorphanol in selegiline-treated dogs. *Veterinary Anaesthesia and Analgesia* 31:129–137.
- Doenicke A, Roizen MF, Hoernecke R, et al. 1997. Haemolysis after etomidate: comparison of propylene glycol and lipid formulations. *British Journal of Anaesthesia* 79(3):386–388.
- Doerning BJ, Brammer DW, Chrisp CE, et al. 1992. Nephrotoxicity of tiletamine in New Zealand white rabbits. *Laboratory Animal Science* 42(3):267–269.
- Doherty TJ, Geiser DR, Rohrbach BW. 1997. Effect of acepromazine and butorphanol on halothane minimum alveolar concentration in ponies. *Equine Veterinary Journal* 29:374–376.
- Driessen B, Zarucco L, Steffey EP, et al. 2002. Serum fluoride concentrations, biochemical and histopathological changes associated with prolonged sevoflurane anaesthesia in horses. *Journal of Veterinary Medicine. A, Physiology, Pathology, Clinical Medicine* 49(7):337–347.
- DuBois RN, Abramson SB, Crofford L, et al. 1998. Cyclooxygenase in biology and disease. *FASEB Journal* 12:1063–1073.
- Dunn AM, Buckley BM. 1986. Non-steroidal anti-inflammatory drugs and the kidney. *British Medical Journal* 293:202–203.
- Dutta NK, Annadurai S, Mazumdar K, et al. 2007. Potential management of resistant microbial infections with a novel non-antibiotic: the anti-inflammatory drug diclofenac sodium. *International Journal of Antimicrobial Agents* 30(3):242–249.
- Eger EI 2nd, Raines DE, Shafer SL, et al. 2008. Is a new paradigm needed to explain how inhaled anesthetics produce immobility? *Anesthesia and Analgesia* 107(3):832–848.
- Egger CM, Duke T, Archer J, et al. 1998. Comparison of plasma fentanyl concentrations by using three transdermal fentanyl patch sizes in dogs. *Veterinary Surgery* 27:159–166.
- Egger CM, Glerum LE, Allen SW, et al. 2003. Plasma fentanyl concentrations in awake cats and cats undergoing anesthesia and ovariohysterectomy using transdermal administration. *Veterinary Anaesthesia and Analgesia* 30:229–236.
- el Bindary EM, Abu el-Nasr LM. 2001. Urodynamic changes following intrathecal administration of morphine and fentanyl to dogs. *Eastern Mediterranean Health Journal* 7:189–196.
- Faura CC, Collins SL, Moore RA, et al. 1998. Systematic review of factors affecting the ratios of morphine and its major metabolites. *Pain* 74:43–53.

- Ferre PJ, Pasloske K, Whittem T, et al. 2006. Plasma pharmacokinetics of alfaxalone in dogs after an intravenous bolus of Alfaxan-CD RTU. *Veterinary Anaesthesia and Analgesia* 33:229–236.
- Fields HL, Emson PC, Leigh BK, et al. 1980. Multiple opiate receptor sites on primary afferent fibres. *Nature* 284:351–353.
- Forman SA, Miller KW. 2016. Mapping general anesthetic sites in heteromeric  $\gamma$ -aminobutyric acid type A receptors reveals a potential for targeting receptor subtypes. *Anesthesia and Analgesia* 123:1263–1273.
- Franks NP. 2006. Molecular targets underlying general anaesthesia. *British Journal of Pharmacology* 147 (Suppl 1):S72–81.
- Franks NP. 2008. General anaesthesia: from molecular targets to neuronal pathways of sleep and arousal. *Nature Reviews. Neuroscience* 9(5):370–386.
- Franks NP, Lieb WR. 1978. Where do general anaesthetics act? *Nature* 274(5669):339–342.
- Franks NP, Lieb WR. 1984. Do general anaesthetics act by competitive binding to specific receptors? *Nature* 310(5978):599–601.
- Franks JN, Boothe HW, Taylor L, et al. 2000. Evaluation of transdermal fentanyl patches for analgesia in cats undergoing onychectomy. *Journal of the American Veterinary Medical Association* 217(7):1013–1020.
- Gellasch KL, Kruse-Elliott KT, Osmond CS, et al. 2002. Comparison of transdermal administration of fentanyl versus intramuscular administration of butorphanol for analgesia after onychectomy in cats. *Journal of the American Veterinary Medical Association* 220:1020–1024.
- Gilberto DB, Motzel SL, Das SR. 2003. Postoperative pain management using fentanyl patches in dogs. *Contemporary Topics in Laboratory Animal Science* 42:21–26.
- Gingerich DA, Baggot JD, Yeary RA. 1975. Pharmacokinetics and dosage of aspirin in cattle. *Journal of the American Veterinary Medical Association* 167(10):945–948.
- Gomez de Segura IA, de la Vibora JB, Aguado D. 2009. Opioid tolerance blunts the reduction in the sevoflurane minimum alveolar concentration produced by remifentanyl in the rat. *Anesthesiology* 110:1133–1138.
- Gorman AL, Elliott KJ, Inturrisi CE. 1997. The d- and l-isomers of methadone bind to the non-competitive site on the N-methyl-D-aspartate (NMDA) receptor in rat forebrain and spinal cord. *Neuroscience Letters* 223:5–8.
- Gozalo-Marcilla M, Luna SPL, Gasthuys F, et al. 2019. Clinical applicability of detomidine and methadone constant rate infusions for surgery in standing horses. *Veterinary Anaesthesia and Analgesia* 46(3):325–334.
- Granhölm M, McKusick BC, Westerholm FC, et al. 2006. Evaluation of the clinical efficacy and safety of dexmedetomidine or medetomidine in cats and their reversal with atipamezole. *Veterinary Anaesthesia and Analgesia* 33:214–223.
- Granhölm M, McKusick BC, Westerholm FC, et al. 2007. Evaluation of the clinical efficacy and safety of intramuscular and intravenous doses of dexmedetomidine and medetomidine in dogs and their reversal with atipamezole. *The Veterinary Record* 160:891–897.
- Grimm JB, Grimm KA, Kneller SK, et al. 2001. The effect of a combination of medetomidine-butorphanol and medetomidine, butorphanol, atropine on glomerular filtration rate in dogs. *Veterinary Radiology and Ultrasound* 42(5):458–462.
- Gutstein HB, Akil H. Opioid analgesics. 2001. In: *Goodman and Gilman's: The Pharmacological Basis of Therapeutics* (JG Harman, LE Limbird, A Goodman Gilman, eds.), p. 569. New York: McGraw-Hill.
- Harrison LM, Kastin AJ, Zadina JE. 1998. Opiate tolerance and dependence: Receptors, G-proteins, and antiopiates. *Peptides* 19:1603–1630.
- Heinonen EH, Myllylä V. 1998. Safety of selegiline (deprenyl) in the treatment of Parkinson's disease. *Drug Safety* 19:11–22.
- Hellyer PW, Bai L, Supon J, et al. 2003. Comparison of opioid and alpha-2 adrenergic receptor binding in horse and dog brain using radioligand autoradiography. *Veterinary Anaesthesia and Analgesia* 30:172–182.
- Hikasa Y, Akiba T, Iino Y, et al. 1992. Central alpha-adrenoceptor subtypes involved in the emetic pathway in cats. *European Journal of Pharmacology* 229(2–3):241–251.
- Hoffman WE, Charbel FT, Edelman G, et al. 1995. Nitrous oxide added to isoflurane increases brain artery blood flow and low frequency brain electrical activity. *Journal of Neurosurgical Anesthesiology* 7(2):82–88.
- Hoke JF, Cunningham F, James MK, et al. 1997. Comparative pharmacokinetics and pharmacodynamics of remifentanyl, its principle metabolite (GR90291) and alfentanil in dogs. *The Journal of Pharmacology and Experimental Therapeutics* 281:226–232.
- Hudetz AG, Mashour GA. 2016. Disconnecting consciousness: is there a common anesthetic end point? *Anesthesia and Analgesia* 123:1228–1240.
- Hudson N, Balsitis M, Everitt S, et al. 1993. Enhanced gastric mucosal leukotriene B4 synthesis in patients taking non-steroidal anti-inflammatory drugs. *Gut* 34:742–747.
- Hunt JR, Grint NJ, Taylor PM, et al. 2013. Sedative and analgesic effects of buprenorphine, combined with either acepromazine or dexmedetomidine, for premedication prior to elective surgery in cats and dogs. *Veterinary Anaesthesia and Analgesia* 40(3):297–307.
- Hussain I, Khan MZ, Khan A, et al. 2008. Toxicological effects of diclofenac in four avian species. *Avian Pathology* 37(3):315–321.

- Ilkiw JE, Pascoe PJ, Fisher LD. 1997. Effect of alfentanil on the minimum alveolar concentration of isoflurane in cats. *American Journal of Veterinary Research* 58:1274–1279.
- Ilkiw JE, Pascoe PJ, Tripp LD. 2002. Effects of morphine, butorphanol, buprenorphine, and U50488H on the minimum alveolar concentration of isoflurane in cats. *American Journal of Veterinary Research* 63(8):1198–1202.
- Inturrisi CE. 2002. Clinical pharmacology of opioids for pain. *The Clinical Journal of Pain* 18:S3–13.
- Jain T, Koley KM, Vadlamudi VP, et al. 2009. Diclofenac-induced biochemical and histopathological changes in white leghorn birds (*Gallus domesticus*). *Indian Journal of Pharmacology* 41(5):237–241.
- Janecka A, Fichna J, Janecki T. 2004. Opioid receptors and their ligands. *Current Topics in Medicinal Chemistry* 4:1–17.
- Jayakumar K, Mohan K, Swamy HD, et al. 2010. Study of nephrotoxic potential of acetaminophen in birds. *Toxicology International* 17(2):86–89.
- Jenkins A, Greenblatt EP, Faulkner HJ, et al. 2001. Evidence for a common binding cavity for three general anesthetics within the GABA(A) receptor. *Journal of Neuroscience* 21(6):RC136.
- Jenkins J, Andreasen A, Trudell JR, et al. 2002. Tryptophan scanning mutagenesis in TM4 of the GABA(A) receptor  $\alpha 1$  subunit: implications for modulation by inhaled anesthetics and ion channel structure. *Neuropharmacology* 43:669–678.
- Jones RM, Detmer M, Hill AB, et al. 1981. Incidence of choledochoduodenal sphincter spasm during fentanyl-supplemented anesthesia. *Anesthesia and Analgesia* 60:638–640.
- Kay-Mugford P, Benn SJ, LaMarre J, et al. 2000. In vitro effects of nonsteroidal anti-inflammatory drugs on cyclooxygenase activity in dogs. *American Journal of Veterinary Research* 61:802–810.
- Kearns KS, Swenson B, Ramsay EC. 2000. Oral induction of anesthesia with droperidol and transmucosal carfentanil citrate in chimpanzees (*Pan troglodytes*). *Journal of Zoo and Wildlife Medicine* 31:185–189.
- Kharasch ED, Schroeder JL, Sheffels P, et al. 2005. Influence of sevoflurane on the metabolism and renal effects of compound A in rats. *Anesthesiology* 103(6):1183–1188.
- Kieffer BL. 1999. Opioids: first lessons from knockout mice. *Trends in Pharmacological Sciences* 20:19–26.
- Kirchner T, Aparicio B, Argentieri DC, et al. 1997a. Effects of tepoxalin, a dual inhibitor of cyclooxygenase/5-lipoxygenase, on events associated with NSAID-induced gastrointestinal inflammation. *Prostaglandins, Leukotrienes, and Essential Fatty Acids* 56:417–423.
- Kirchner T, Argentieri DC, Barbone AG, et al. 1997b. Evaluation of the antiinflammatory activity of a dual cyclooxygenase-2 selective/5-lipoxygenase inhibitor, RWJ 63556, in a canine model of inflammation. *The Journal of Pharmacology and Experimental Therapeutics* 282:1094–1101.
- Kirkby Shaw K, Rausch-Derra LC, Rhodes L. 2015. Grapiprant: an EP4 prostaglandin receptor antagonist and novel therapy for pain and inflammation. *Veterinary Medicine and Science* 2(1):3–9.
- Ko JC, Lange DN, Mandsager RE, et al. 2000. Effects of butorphanol and carprofen on the minimal alveolar concentration of isoflurane in dogs. *Journal of the American Veterinary Medical Association* 217(7):1025–1028.
- Kohrs R, Durieux ME. 1998. Ketamine: teaching an old drug new tricks. *Anesthesia and Analgesia* 87(5):1186–1193.
- Krasowski MD, Harrison NL. 2000. The actions of ether, alcohol and alkane general anaesthetics on GABA(A) and glycine receptors and the effects of TM2 and TM3 mutations. *British Journal of Pharmacology* 129:731–743.
- Kuipers PW, Kamphuis ET, van Venrooij GE, et al. 2004. Intrathecal opioids and lower urinary tract function: a urodynamic evaluation. *Anesthesiology* 100:1497–1503.
- Kuusela E, Raekallio M, Anttila M, et al. 2000. Clinical effects and pharmacokinetics of medetomidine and its enantiomers in dogs. *Journal of Veterinary Pharmacology and Therapeutics* 23:15–20.
- Laredo FG, Sanchez-Valverde MA, Cantalapiedra AG, et al. 1998. Efficacy of the Komesaroff anaesthetic machine for delivering isoflurane to dogs. *The Veterinary Record* 43(16):437–440.
- Lascelles BD, Robertson SA. 2004. Antinociceptive effects of hydromorphone, butorphanol, or the combination in cats. *Journal of Veterinary Internal Medicine* 18:190–195.
- Lee HK, Wang SC. 1975. Mechanism of morphine-induced miosis in the dog. *The Journal of Pharmacology and Experimental Therapeutics* 192:415–431.
- Lee DD, Papich MG, Hardie EM. 2000. Comparison of pharmacokinetics of fentanyl after intravenous and transdermal administration in cats. *American Journal of Veterinary Research* 61:672–677.
- Lee M, Silverman S, Hansen H, et al. 2011. A comprehensive review of opioid-induced hyperalgesia. *Pain Physician* 14:145–161.
- Lerman J, Robinson S, Willis MM, et al. 1983. Anesthetic requirements for halothane in young children 0–1 month and 1–6 months of age. *Anesthesiology* 59(5):421–424.
- Lerman J, Sikich N, Kleinman S, et al. 1994. The pharmacology of sevoflurane in infants and children. *Anesthesiology* 80(4):814–824.
- Li D, Hambrecht-Wiedbusch VS, Mashour GA. 2017. Accelerated recovery of consciousness after general anesthesia is associated with increased functional brain

- connectivity in the high-gamma bandwidth. *Frontiers in Systems Neuroscience* 11:16.
- Livingston A. 2000. Mechanism of action of nonsteroidal anti-inflammatory drugs. *The Veterinary Clinics of North America. Small Animal Practice* 30:773–781.
- Lobetti RG, Joubert KE. 2000. Effect of administration of nonsteroidal anti-inflammatory drugs before surgery on renal function in clinically normal dogs. *American Journal of Veterinary Research* 61:1501–1507.
- MacAllister CG, Morgan SJ, Borne AT, et al. 1993. Comparison of adverse effects of phenylbutazone, flunixin meglumine, and ketoprofen in horses. *Journal of the American Veterinary Medical Association* 202:71–77.
- Machin KL, Caulkett NA. 1998. Cardiopulmonary effects of propofol and a medetomidine-midazolam-ketamine combination in mallard ducks. *American Journal of Veterinary Research* 59(5):598–602.
- Mackiewicz AL, Salyards GW, Knych HK, et al. 2019. Pharmacokinetics of a long-lasting, highly concentrated buprenorphine solution after subcutaneous administration in Rhesus Macaques (*Macaca mulatta*). *Journal of the American Association for Laboratory Animal Science* 58(4):501–509.
- Malmberg AB, Yaksh TL. 1992. Antinociceptive actions of spinal nonsteroidal anti-inflammatory agents on the formalin test in the rat. *The Journal of Pharmacology and Experimental Therapeutics* 263:136–146.
- Mama KR, Steffey EP, Withrow SJ. 2000. Use of orally administered carfentanil prior to isoflurane-induced anesthesia in a Kodiak brown bear. *Journal of the American Veterinary Medical Association* 217:546–549.
- Mashour GA, Hudetz AG. 2017. Bottom-up and top-down mechanisms of general anesthetics modulate different dimensions of consciousness. *Frontiers in Neural Circuits* 11:44.
- Mathews KA, Pettifer G, Foster RF. 1999. A comparison of the safety and efficacy of meloxicam to ketoprofen and butorphanol for control of post-operative pain associated with soft tissue surgery in dogs. Proceedings of the Symposium on Recent Advances in Non-Steroidal Anti-Inflammatory Therapy in Small Animals, p. 67. Paris.
- Mathies T, Mauldin RE. 2020. Lethal methemoglobinemia in the invasive brown treesnake after acetaminophen ingestion. *Scientific Reports* 10(1):845.
- Matthews NS, Lindsay SL. 1990. Effect of low-dose butorphanol on halothane minimum alveolar concentration in ponies. *Equine Veterinary Journal* 22:325–327.
- Maxwell LK, Thomasy SM, Slovis N, et al. 2003. Pharmacokinetics of fentanyl following intravenous and transdermal administration in horses. *Equine Veterinary Journal* 35:484–490.
- Maze M, Tranquilli W. 1991. Alpha-2 adrenoceptor agonists: defining the role in clinical anesthesia. *Anesthesiology* 74(3):581–605.
- McCann ME, Andersen DR, Zhang D, et al. 2004. In vitro effects and in vivo efficacy of a novel cyclooxygenase-2 inhibitor in dogs with experimentally induced synovitis. *American Journal of Veterinary Research* 65:503–512.
- McCormack K. 1994. Non-steroidal anti-inflammatory drugs and spinal nociceptive processing. *Pain* 59:9–43.
- McKellar QA, Pearson T, Bogan JA. 1990. Pharmacokinetics, tolerance and serum thromboxane inhibition of carprofen in the dog. *The Journal of Small Animal Practice* 31:443–448.
- McKellar QA, Delatour P, Lees P. 1994a. Stereospecific pharmacodynamics and pharmacokinetics of carprofen in the dog. *Journal of Veterinary Pharmacology and Therapeutics* 17:447–454.
- McKellar QA, Lees P, Gettinby G. 1994b. Pharmacodynamics of tolafenamic acid in dogs. Evaluation of dose response relationships. *European Journal of Pharmacology* 253:191–200.
- Mendes GM, Selmi AL. 2004. Use of a combination of propofol and fentanyl, alfentanil, or sufentanil for total intravenous anesthesia in cats. *Journal of the American Veterinary Med Association* 223:1608–1613.
- Mercadante S, Arcuri E. 2004. Opioids and renal function. *Journal of Pain* 5:2–19.
- Merkel G, Eger E. 1963. A comparative study of halothane and halopropane anesthesia: including method for determining equipotency. *Anesthesiology* 23:346–357.
- Meyer HH. 1901. Zur Theorie der Alkoholnarkose III. Der Einfluss wechselnder Temperatur auf Wirkungs-stärke und Teilungs Koeffizient der Narkotika. *Archiv für experimentelle Pathologie und Pharmakologie* 154:338–346.
- Michelsen LG, Salmenpera M, Hug CC Jr., et al. 1996. Anesthetic potency of remifentanyl in dogs. *Anesthesiology* 84:865–872.
- Mielck F, Stephan H, Buhre W, et al. 1998. Effects of 1 MAC desflurane on cerebral metabolism, blood flow and carbon dioxide reactivity in humans. *British Journal of Anaesthesia* 81(2):155–160.
- Mielck F, Stephan H, Weyland A, et al. 1999. Effects of one minimum alveolar anesthetic concentration sevoflurane on cerebral metabolism, blood flow, and CO<sub>2</sub> reactivity in cardiac patients. *Anesthesia and Analgesia* 89(2):364–369.
- Mihic SJ, Ye Q, Wick MJ, et al. 1997. Sites of alcohol and volatile anaesthetic action on GABA(A) and glycine receptors. *Nature* 389(6649):385–389.
- Miller BF, Muller LI, Storms TN, et al. 2003. A comparison of carfentanil/xylazine and telazol (R)/xylazine for

- immobilization of white-tailed deer. *Journal of Wildlife Diseases* 39:851–858.
- Millis DL, Weigel JP, Moyers T, et al. 2002. Effect of deracoxib, a new COX-2 inhibitor, on the prevention of lameness induced by chemical synovitis in dogs. *Veterinary Therapeutics* 3:453–464.
- Moon PF. 1994. Acute toxicosis in two dogs associated with etomidate-propylene glycol infusion. *Laboratory Animal Science* 44(6):590–594.
- Moon PF. 1997. Cortisol suppression in cats after induction of anesthesia with etomidate, compared with ketamine-diazepam combination. *American Journal of Veterinary Research* 58(8):868–871.
- Moran TD, Abdulla FA, Smith PA. 2000. Cellular neurophysiological actions of nociceptin/orphanin FQ. *Peptides* 21:969–976.
- Moreau M, Daminet S, Martel-Pelletier J, et al. 2005. Superiority of the gastroduodenal safety profile of licofelone over rofecoxib, a COX-2 selective inhibitor, in dogs. *Journal of Veterinary Pharmacology and Therapeutics* 28:81–86.
- Mortenson J, Bechert U. 2001. Carfentanil citrate used as an oral anesthetic agent for brown bears (*Ursus arctos*). *Journal of Zoo and Wildlife Medicine* 32:217–221.
- Mosely CI, Mosely CA. 2015. Comparative anesthesia and analgesia of reptiles, amphibians and fishes. In: *Veterinary Anesthesia and Analgesia: The Fifth Edition of Lumb and Jones* (KA Grimm, KA Lamont, WJ Tranquilli, SA Greene, SA Robertson, eds.). Ames: Wiley Blackwell.
- Muir WW 3rd, Gadawski JE. 1998. Respiratory depression and apnea induced by propofol in dogs. *American Journal of Veterinary Research* 59(2):157–161.
- Muir W, Lerche P, Wiese A, et al. 2008. Cardiorespiratory and anesthetic effects of clinical and supraclinical doses of alfaxalone in dogs. *Veterinary Anaesthesia and Analgesia* 35:451–462.
- Muir W, Lerche P, Wiese A, et al. 2009. The cardiorespiratory and anesthetic effects of clinical and supraclinical doses of alfaxalone in cats. *Veterinary Anaesthesia and Analgesia* 36:42–54.
- Naidoo V, Wolter K, Cuthbert R, et al. 2009. Veterinary diclofenac threatens Africa's endangered vulture species. *Regulatory Toxicology and Pharmacology* 53(3):205–208.
- Natalini CC, Robinson EP. 2000. Evaluation of the analgesic effects of epidurally administered morphine, alfentanil, butorphanol, tramadol, and U50488H in horses. *American Journal of Veterinary Research* 61:1579–1586.
- Neuberger J. 1998. Halothane hepatitis. *European Journal of Gastroenterology & Hepatology* 10(8):631–633.
- Niedfeldt RL, Robertson SA. 2006. Postanesthetic hyperthermia in cats: a retrospective comparison between hydromorphone and buprenorphine. *Veterinary Anaesthesia and Analgesia* 33(6):381–389.
- Nishikawa K, Harrison NL. 2003. The actions of sevoflurane and desflurane on the g-aminobutyric acid receptor type A – effects of TM2 mutations in the alpha and beta subunits. *Anesthesiology* 99:678–684.
- Novartis Animal Health USA. 2004. Deracoxib Package Insert.
- Oaks JL, Gilbert M, Virani MZ, et al. 2004. Diclofenac residues as the cause of vulture population decline in Pakistan. *Nature* 427(6975):630–633.
- Osman TE, Nicholson T. 1991. Alpha-2 adrenoreceptors mediate clonidine-induced hypoinsulinaemia in sheep. *Journal of Veterinary Pharmacology and Therapeutics* 14(3):293–299.
- Overton CE. 1901. Studien über die Narkose zugleich ein Beitrag zur allgemeinen Pharmakologie. Jena: Gustav Fischer.
- Page SW, Maddison JE. 2002. Principles of clinical pharmacology. In: *Small Animal Clinical Pharmacology* (J Maddison, S Page, D Church, eds.). London: W.B. Saunders.
- Pascoe PJ, Taylor PM. 2003. Effects of dopamine antagonists on alfentanil-induced locomotor activity in horses. *Veterinary Anaesthesia and Analgesia* 30:165–171.
- Pascoe PJ, Black WD, Claxton JM, et al. 1991. The pharmacokinetics and locomotor activity of alfentanil in the horse. *Journal of Veterinary Pharmacology and Therapeutics* 14:317–325.
- Pascoe PJ, Steffey EP, Black WD, et al. 1993. Evaluation of the effect of alfentanil on the minimum alveolar concentration of halothane in horses. *American Journal of Veterinary Research* 54:1327–1332.
- Pascoe PJ, Ilkiw JE, Fisher LD. 1997. Cardiovascular effects of equipotent isoflurane and alfentanil/isoflurane minimum alveolar concentration multiple in cats. *American Journal of Veterinary Research* 58:1267–1273.
- Patel AJ, Honoré E, Lesage F, et al. 1999. Inhalational anesthetics activate two-pore-domain background K<sup>+</sup> channels. *Nature Neuroscience* 2:422–426.
- Petersson I, Nilsson G, Hansson BG, et al. 1987. Water intoxication associated with non-steroidal anti-inflammatory drug therapy. *Acta Medica Scandinavica* 221:221–223.
- Pettifer GR, Hosgood G. 2003. The effect of rectal temperature on perianesthetic serum concentrations of transdermally administered fentanyl in cats anesthetized with isoflurane. *American Journal of Veterinary Research* 64:1557–1561.
- Pibarot P, Dupuis J, Grisneaux E, et al. 1997. Comparison of ketoprofen, oxymorphone hydrochloride, and butorphanol in the treatment of postoperative pain in

- dogs. *Journal of the American Veterinary Medical Association* 211:438–444.
- Pollock CG, Ramsay EC. 2003. Serial immobilization of a Brazilian tapir (*Tapirus terrestris*) with oral detomidine and oral carfentanil. *Journal of Zoo and Wildlife Medicine* 34:408–410.
- Posner LP, Gleed RD, Erb HN, et al. 2007. Post-anesthetic hyperthermia in cats. *Veterinary Anaesthesia and Analgesia* 34(1):40–47.
- Posner LP, Pavuk AA, Rokshar JL, et al. 2010. Effects of opioids and anesthetic drugs on body temperature in cats. *Veterinary Anaesthesia and Analgesia* 37(1):35–43.
- Poulsen Nautrep B, Justus C. 1999. Effects of some veterinary NSAIDs on ex vivo thromboxane production and in vivo urine output in the dog. Proceedings of the Symposium on Recent Advances in Non-Steroidal Anti-Inflammatory Therapy in Small Animals, p. 25. Paris.
- Priano LL, Vatner SF. 1981. Generalized cardiovascular and regional hemodynamic effects of meperidine in conscious dogs. *Anesthesia and Analgesia* 60:649–654.
- Pypendop BH, Verstegen JP. 1998. Hemodynamic effects of medetomidine in the dog: a dose titration study. *Veterinary Surgery* 27(6):612–622.
- Quandt JE, Raffae MR, Robinson EP. 1994. Butorphanol does not reduce the minimum alveolar concentration of halothane in dogs. *Veterinary Surgery* 23:156–159.
- Radnay PA, Duncalf D, Novakovic M, et al. 1984. Common bile duct pressure changes after fentanyl, morphine, meperidine, butorphanol, and naloxone. *Anesthesia and Analgesia* 63:441–444.
- Rainsford KD. 1992. Mechanisms of NSAID-induced ulcerogenesis: structural properties of drugs, focus on the microvascular factors, and novel approaches for gastrointestinal protection. *Acta Physiologica Hungarica* 80:23–38.
- Ramdohr S, Bornemann H, Plotz J, et al. 2001. Immobilization of free-ranging adult male southern elephant seals with Immobilon (TM) (etorphine/acepromazine) and ketamine. *South African Journal of Wildlife Research* 31:135–140.
- Rausch-Derra L, Huebner M, Wofford J, et al. 2016. A prospective, randomized, masked, placebo-controlled multisite clinical study of grapiprant, an EP4 prostaglandin receptor antagonist (PRA), in dogs with osteoarthritis. *Journal of Veterinary Internal Medicine* 30(3):756–763.
- Reed S. 2002. Nonsteroidal anti-inflammatory drug-induced duodenal ulceration and perforation in a mature rottweiler. *The Canadian Veterinary Journal* 43:971–972.
- Ricciotti E, FitzGerald GA. 2011. Prostaglandins and inflammation. *Arteriosclerosis, Thrombosis, and Vascular Biology* 31:986–1000.
- Ricketts AP, Lundy KM, Seibel SB. 1998. Evaluation of selective inhibition of canine cyclooxygenase 1 and 2 by carprofen and other nonsteroidal anti-inflammatory drugs. *American Journal of Veterinary Research* 59:1441–1446.
- Ripamonti C, Dickerson ED. 2001. Strategies for the treatment of cancer pain in the new millennium. *Drugs* 61:955–977.
- Roald OK, Forsman M, Heier MS, et al. 1991. Cerebral effects of nitrous oxide when added to low and high concentrations of isoflurane in the dog. *Anesthesia and Analgesia* 72(1):75–79.
- Robertson SA, Taylor PM, Lascelles BDX, et al. 2003a. Changes in thermal threshold response in eight cats after administration of buprenorphine, butorphanol and morphine. *The Veterinary Record* 153:462–465.
- Robertson SA, Taylor PM, Sear JW. 2003b. Systemic uptake of buprenorphine by cats after oral mucosal administration. *The Veterinary Record* 152:675–678.
- Robinson TM, Kruse-Elliott KT, Markel MD, et al. 1999. A comparison of transdermal fentanyl versus epidural morphine for analgesia in dogs undergoing major orthopedic surgery. *Journal of the American Animal Hospital Association* 35:95–100.
- Robinson MA, Guan F, McDonnell S, et al. 2015. Pharmacokinetics and pharmacodynamics of dermorphin in the horse. *Journal of Veterinary Pharmacology and Therapeutics* 38(4):321–329.
- Roffe TJ, Coffin K, Berger J. 2001. Survival and immobilizing moose with carfentanil and xylazine. *Wildlife Society Bulletin* 29:1140–1146.
- Ruiz-López P, Navarrete-Calvo R, Morgaz J, et al. 2020. Determination of acute tolerance and hyperalgesia to remifentanyl constant rate infusion in dogs undergoing sevoflurane anaesthesia. *Veterinary Anaesthesia and Analgesia* 47(2):183–190.
- Saleh N, Aoki M, Shimada T, et al. 2005. Renal effects of medetomidine in isoflurane-anesthetized dogs with special reference to its diuretic action. *The Journal of Veterinary Medical Science* 67(5):461–465.
- Sanhouri AA, Jones RS, Dobson H. 1992. Effects of xylazine on the stress response to transport in male goats. *The British Veterinary Journal* 148(2):119–128.
- Santonocito C, Noto A, Crimi C, et al. 2018. Remifentanyl-induced postoperative hyperalgesia: current perspectives on mechanisms and therapeutic strategies. *Local and Regional Anesthesia* 11:15–23.
- Sawyer DC, Rech RH, Durham RA, et al. 1991. Dose response to butorphanol administered subcutaneously to increase visceral nociceptive threshold in dogs. *American Journal of Veterinary Research* 52:1826–1830.
- Sellon DC, Monroe VL, Roberts MC, et al. 2001. Pharmacokinetics and adverse effects of butorphanol

- administered by single intravenous injection or continuous intravenous infusion in horses. *American Journal of Veterinary Research* 62:183–189.
- Serhan CN, Chiang N. 2004. Novel endogenous small molecules as the checkpoint controllers in inflammation and resolution: entree for resolomics. *Rheumatic Diseases Clinics of North America* 30:69–95.
- Shah MD, Yates D, Hunt J, et al. 2018. A comparison between methadone and buprenorphine for perioperative analgesia in dogs undergoing ovariohysterectomy. *The Journal of Small Animal Practice* 59(9):539–546.
- Sheffels P, Schroeder JL, Altuntas TG, et al. 2004. Role of cytochrome P4503A in cysteine S-conjugates sulfoxidation and the nephrotoxicity of the sevoflurane degradation product fluoromethyl-2,2-difluoro-1-(trifluoromethyl)vinyl ether (compound A) in rats. *Chemical Research in Toxicology* 17(9):1177–1189.
- Short CE. 1991. Effects of anticholinergic treatment on the cardiac and respiratory systems in dogs sedated with medetomidine. *The Veterinary Record* 129(14):310–313.
- Silverstein FE, Faich G, Goldstein JL, et al. 2000. Gastrointestinal toxicity with celecoxib vs nonsteroidal anti-inflammatory drugs for osteoarthritis and rheumatoid arthritis: the CLASS study: A randomized controlled trial. Celecoxib Long-term Arthritis Safety Study. *JAMA: The Journal of the American Medical Association* 284:1247–1255.
- Sinclair, M, Mealey KL, Mathews NS, et al. 2003. The pharmacokinetics of meloxicam in horses. Proceedings of the 8th World Congress of Veterinary Anesthesia, p. 112. Knoxville, TN.
- Sleigh J, Warnaby C, Tracey I. 2018. General anaesthesia as fragmentation of selfhood: insights from electroencephalography and neuroimaging. *British Journal of Anaesthesia* 121:233–240.
- Slingsby LS, Bortolami E, Murrell JC. 2015. Methadone in combination with medetomidine as premedication prior to ovariohysterectomy and castration in the cat. *Journal of Feline Medicine and Surgery* 17(10):864–872.
- Smith MT. 2000. Neuroexcitatory effects of morphine and hydromorphone: evidence implicating the 3-glucuronide metabolites. *Clinical and Experimental Pharmacology & Physiology* 27:524–528.
- Smith AP, Lee NM. 2003. Opioid receptor interactions: Local and nonlocal, symmetric and asymmetric, physical and functional. *Life Sciences* 73:1873–1893.
- Smith PA, Moran TD. 2001. The nociceptin receptor as a potential target in drug design. *Drug News & Perspectives* 14:335–345.
- Smith LJ, Yu JKA, Bjorling DE, et al. 2001. Effects of hydromorphone or oxymorphone, with or without acepromazine, on preanesthetic sedation, physiologic values, and histamine release in dogs. *Journal of the American Veterinary Medical Association* 218:1101–1105.
- Snow DH, Douglas TA, Thompson H, et al. 1981. Phenylbutazone toxicosis in equidae: a biochemical and pathophysiological study. *American Journal of Veterinary Research* 42:1754–1759.
- Sporer KA. 1995. The serotonin syndrome. Implicated drugs, pathophysiology and management. *Drug Safety* 13:94–104.
- Steffey EP, Mama KM, Brosnan RJ. 2015. Inhalation anesthetics. In: *Veterinary Anesthesia and Analgesia: The Fifth Edition of Lumb and Jones* (KA Grimm, KA Lamont, WJ Tranquilli, SA Greene, SA Robertson, eds.) Ames: Wiley Blackwell.
- Stein C. 1993. Peripheral mechanisms of opioid analgesia. *Anesthesia and Analgesia* 76:182–191.
- Stein C. 1995. The control of pain in peripheral tissue by opioids. *New England Journal of Medicine* 332:1685–1690.
- Stein C, Yassouridis A. 1997. Peripheral morphine analgesia. *Pain* 71:119–121.
- Stein C, Hassan AH, Lehrberger K, et al. 1993. Local analgesic effect of endogenous opioid peptides. *Lancet* 342:321–324.
- Stein C, Pfluger M, Yassouridis A, et al. 1996. No tolerance to peripheral morphine analgesia in presence of opioid expression in inflamed synovia. *The Journal of Clinical Investigation* 98:793–799.
- Stein C, Machelska H, Schafer M. 2001. Peripheral analgesic and antiinflammatory effects of opioids. *Zeitschrift fur Rheumatologie* 60:416–424.
- Stein C, Schafer M, Machelska H. 2003. Attacking pain at its source: new perspectives on opioids. *Nature Medicine* 9:1003–1008.
- Stephan DD, Vestre WA, Stiles J, et al. 2003. Changes in intraocular pressure and pupil size following intramuscular administration of hydromorphone hydrochloride and acepromazine in clinically normal dogs. *Veterinary Ophthalmology* 6:73–76.
- Stoelting RK. 1999. Opioid agonists and antagonists. In: *Pharmacology and Physiology in Anesthetic Practice* (RK Stoelting, ed.), p. 77. Philadelphia: Lippincott Williams & Wilkins.
- Stone PA, Macintyre PE, Jarvis DA. 1993. Norpethidine toxicity and patient controlled analgesia. *British Journal of Anaesthesia* 71:738–740.
- Taggart MA, Senacha KR, Green RE, et al. 2007. Diclofenac residues in carcasses of domestic ungulates available to vultures in India. *Environment International* 33(6):759–765.
- Takada K, Tonner PH, Maze M. 1999. Meperidine functions as an alpha2B adrenoceptor agonist. *Anesthesiology* 91:U363.

- Takada K, Clark DJ, Davies MF, et al. 2002. Meperidine exerts agonist activity at the alpha(2B)-adrenoceptor subtype. *Anesthesiology* 96:1420–1426.
- Taylor RH, Lerman J. 1991. Minimum alveolar concentration of desflurane and hemodynamic responses in neonates, infants, and children. *Anesthesiology* 75(6):975–979.
- Taylor PM, Robertson SA, Dixon MJ, et al. 2001. Morphine, pethidine and buprenorphine disposition in the cat. *Journal of Veterinary Pharmacology and Therapeutics* 24:391–398.
- Thompson JR, Kersting KW, Hsu WH. 1991. Antagonistic effect of atipamezole on xylazine-induced sedation, bradycardia, and ruminal atony in calves. *American Journal of Veterinary Research* 52(8):1265–1268.
- Tissot TA. 2003. Probable meperidine-induced serotonin syndrome in a patient with a history of fluoxetine use. *Anesthesiology* 98:1511–1512.
- Trimble T, Bhalla RJ, Leece EA. 2018. Comparison of sedation in dogs: methadone or butorphanol in combination with dexmedetomidine intravenously. *Veterinary Anaesthesia and Analgesia* 45(5):597–603.
- Uhrig L, Dehaene S, Jarraya B. 2014. Cerebral mechanisms of general anesthesia. *Annales Francaises d'Anesthesie et de Reanimation* 33(2):72–82.
- US Food and Drug Administration. 2011. Center for Veterinary Medicine Freedom of Information Summary for Firocoxib. www.FDA.gov.
- Van de Wiele B, Rubinstein E, Peacock W, et al. 1995. Propylene glycol toxicity caused by prolonged infusion of etomidate. *Journal of Neurosurgical Anesthesiology* 1995;7(4):259–62.
- Vane JR. 1971. Inhibition of prostaglandin synthesis as a mechanism of action for aspirin-like drugs. *Nature: New Biology* 231:232–235.
- Vane JR, Botting RM. 1995. New insights into the mode of action of anti-inflammatory drugs. *Inflammation Research* 44:1–10.
- Vane JR, Botting RM. 2003. The mechanism of action of aspirin. *Thrombosis Research* 110:255–258.
- Veng-Pedersen P, Wilhelm JA, Zakszewski TB, et al. 1995. Duration of opioid antagonism by nalmefene and naloxone in the dog: an integrated pharmacokinetic/pharmacodynamic comparison. *Journal of Pharmaceutical Sciences* 84:1101–1106.
- Verbruggen AM, Akkerdaas LC, Hellebrekers LJ, et al. 2000. The effect of intravenous medetomidine on pupil size and intraocular pressure in normotensive dogs. *The Veterinary Quarterly* 22(3):179–180.
- Wagner LE, Eaton M, Sabnis SS, et al. 1999. Meperidine and lidocaine block of recombinant voltage-dependent Na<sup>+</sup> channels: evidence that meperidine is a local anesthetic. *Anesthesiology* 91:1481–1490.
- Wallenstein MC, Wang SC. 1979. Mechanism of morphine-induced mydriasis in the cat. *The American Journal of Physiology* 236:R292–R296.
- Warne LN, Beths T, Holm M, Bauquier SH. 2013. Comparison of perioperative analgesic efficacy between methadone and butorphanol in cats. *Journal of the American Veterinary Medical Association* 243(6):844–850.
- Waterman AE. 1983. Influence of premedication with xylazine on the distribution and metabolism of intramuscularly administered ketamine in cats. *Research in Veterinary Science* 35(3):285–290.
- Welsh EM, Baxter P, Nolan AM. 1992. Pharmacokinetics of carprofen administered intravenously to sheep. *Research in Veterinary Science* 53:264–266.
- West PM, Fernandez C. 2003. Safety of COX-2 inhibitors in asthma patients with aspirin hypersensitivity. *The Annals of Pharmacotherapy* 37:1497–1501.
- White DM, Mair AR, Martinez-Taboada, F. 2017. Opioid-free anaesthesia in three dogs. *Open Veterinary Journal* 7(2):104–110.
- Whittem T, Pasloske KS, Heit MC, Ranasinghe MG. 2008. The pharmacokinetics and pharmacodynamics of alfaxalone in cats after single and multiple intravenous administration of Alfaxan at clinical and supraclinical doses. *Journal of Veterinary Pharmacology and Therapeutics* 31:571–579.
- Wilkinson AC, Thomas ML, Morse BC. 2001. Evaluation of a transdermal fentanyl system in yucatan miniature pigs. *Contemporary Topics in Laboratory Animal Science* 40:12–16.
- Wolff M, Olschewski A, Vogel W, et al. 2004. Meperidine suppresses the excitability of spinal dorsal horn neurons. *Anesthesiology* 100:947–955.
- Yaksh TL. 1998. Pharmacology and mechanisms of opioid analgesic activity. In: *Anesthesia: Biologic Foundations* (TL Yaksh, C Lynch III, WM Zapol, eds.), p. 921. Philadelphia: Lippincott-Raven.
- Yaksh TL, Dirig DM, Malmberg AB. 1998. Mechanism of action of nonsteroidal anti-inflammatory drugs. *Cancer Investigation* 16:509–527.
- Yeh SY, Krebs HA, Changchit A. 1981. Urinary excretion of meperidine and its metabolites. *Journal of Pharmaceutical Sciences* 70:867–870.
- Yu EH, Tran DH, Lam SW, et al. 2016. Remifentanyl tolerance and hyperalgesia: short-term gain, long-term pain? *Anaesthesia* 71:1347–1362.
- Zadina JE, Martin-Schild S, Gerall AA, et al. 1990. Endomorphins: Novel endogenous mu-opiate receptor agonists in regions of high mu-opiate receptor density. *Neuropeptides: Structure and Function in Biology and Behavior* 897:136–144.

